arXiv:1401.0493v1 [math.NT] 2 Jan 2014

Congruences for $q^{[p/8]} \pmod p$ II

ZHI-HONG SUN

School of Mathematical Sciences, Huaiyin Normal University,
Huaian, Jiangsu 223001, PR China
E-mail: zhihongsun@yahoo.com
Homepage: http://www.hytc.edu.cn/xsjl/szh

ABSTRACT. Let $\mathbb{Z}$ be the set of integers, and let p be a prime of the form $4k+1$. Suppose $q \in \mathbb{Z}$, $2 \nmid q$, $p \nmid q$, $p = c^2+d^2$, $c, d \in \mathbb{Z}$ and $c \equiv 1 \pmod 4$. In this paper we continue to discuss congruences for $q^{[p/8]} \pmod p$ and present new reciprocity laws, but we assume $4p = x^2+qy^2$ or $p = x^2+2qy^2$, where $[\cdot]$ is the greatest integer function and $x, y \in \mathbb{Z}$.

MSC: Primary 11A15, Secondary 11A07, 11E25
Keywords: Reciprocity law; octic residue; congruence; quartic Jacobi symbol

1. Introduction.

Let $\mathbb{Z}$ be the set of integers, $i = \sqrt{-1}$ and $\mathbb{Z}[i] = \{a+bi \mid a, b \in \mathbb{Z}\}$. For any positive odd number m and $a \in \mathbb{Z}$ let $(\frac{a}{m})$ be the (quadratic) Jacobi symbol. (We also assume $(\frac{a}{1}) = 1$.) For our convenience we also define $(\frac{a}{-m}) = (\frac{a}{m})$. Then for any two odd numbers m and n with $m > 0$ or $n > 0$ we have the following general quadratic reciprocity law: $(\frac{m}{n}) = (-1)^{\frac{m-1}{2}\cdot\frac{n-1}{2}}(\frac{n}{m})$.

For $a, b, c, d \in \mathbb{Z}$ with $2 \nmid c$ and $2 \mid d$, one can define the quartic Jacobi symbol $\left(\frac{a+bi}{c+di}\right)_4$ as in [S1,S2,S4]. From [IR] we know that $\overline{\left(\frac{a+bi}{c+di}\right)_4} = \left(\frac{a-bi}{c-di}\right)_4 = \left(\frac{a+bi}{c+di}\right)_4^{-1}$, where $\bar{x}$ means the complex conjugate of x. In Section 2 we list main properties of the quartic Jacobi symbol. See also [IR], [BEW] and [S4]. For the history of quartic reciprocity laws, see [Lem].

Let p be a prime of the form $4k+1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2+d^2 = x^2+qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume that $(c, x+d) = 1$ or $(d_0, x+c) = 1$, where (m, n) is the greatest common divisor of m and n. In [S5], using the quartic reciprocity law the author deduced some congruences for $q^{[p/8]} \pmod p$ in terms of c, d, x and y, where $[\cdot]$ is the greatest integer function. For example, if p and q are primes of the form $8k+1$, $p = c^2+d^2 = x^2+qy^2$, $q = a^2+b^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$, then

$$q^{\frac{p-1}{8}} \equiv (-1)^{\frac{d}{4}+\frac{xy}{4}}\left(\frac{d}{c}\right)^m \pmod p \iff \left(\frac{ac+bd}{ac-bd}\right)^{\frac{q-1}{8}} \equiv \left(\frac{b}{a}\right)^m \pmod q.$$

The author is supported by the National Natural Sciences Foundation of China (No. 11371163).

If $p\equiv 1,9,25\pmod{56}$ is a prime and so $p=c^2+d^2=x^2+7y^2$ for some $c,d,x,y\in\mathbb{Z}$, by [S5, Corollary 4.2] we have

$$7^{\frac{p-1}{8}}\equiv\begin{cases}-(-1)^{\frac{y}{4}}\pmod p & \text{if } 7\mid c,\\ (-1)^{\frac{y}{4}}\pmod p & \text{if } 7\mid d,\\ \pm(-1)^{\frac{y}{4}}\frac{c}{d}\pmod p & \text{if } c\equiv\pm d\pmod 7.\end{cases}$$

Let p be a prime of the form $4k+1$, $q\in\mathbb{Z}$, $2\nmid q$ and $p\nmid q$. Suppose that $p=c^2+d^2$ with $c,d,x,y\in\mathbb{Z}$ and $c\equiv 1\pmod 4$. In this paper we continue to discuss congruences for $q^{[p/8]}\pmod p$, but we assume $4p=x^2+qy^2$ or $p=x^2+2qy^2$. Here are some typical results:

$\star$ Let p be a prime of the form $8k+1$ and $(\frac{-11}{p})=1$, $p=c^2+d^2$, $4p=x^2+11y^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1\pmod 4$ and $d=2^rd_0(2\nmid d_0)$. Assume that $(c,x+2d)=1$ or $(d_0,x+2c)=1$. Then

$$(-11)^{\frac{p-1}{8}}\equiv\begin{cases}\pm(-1)^{[\frac{x}{4}]}\pmod p & \text{if } 2\nmid x \text{ and } x\equiv\pm 4c,\pm 9c\pmod{11},\\ \pm(-1)^{[\frac{x}{4}]}\frac{d}{c}\pmod p & \text{if } 2\nmid x \text{ and } x\equiv\pm 4d,\pm 9d\pmod{11},\\ \mp(-1)^{[\frac{x}{8}]+\frac{y}{8}}\pmod p & \text{if } 2\mid x \text{ and } x\equiv\pm 4c,\pm 9c\pmod{11},\\ \mp(-1)^{[\frac{x}{8}]+\frac{y}{8}}\frac{d}{c}\pmod p & \text{if } 2\mid x \text{ and } x\equiv\pm 4d,\pm 9d\pmod{11}.\end{cases}$$

$\star$ Let p and q be primes such that $p\equiv 1\pmod 8$, $q\equiv 7\pmod 8$, $p=c^2+d^2=x^2+2qy^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1\pmod 4$, $d=2^rd_0$ and $d_0\equiv 1\pmod 4$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then

$$(-q)^{\frac{p-1}{8}}\equiv\Big(\frac{d}{c}\Big)^m\pmod p\iff\Big(\frac{c-di}{c+di}\Big)^{\frac{q+1}{8}}\equiv i^m\pmod q.$$

$\star$ Let p be a prime of the form $8k+1$ and $a\in\mathbb{Z}$ with $2\nmid a$. Suppose that $p=c^2+d^2=x^2+(a^2+1)y^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1\pmod 4$, $d=2^rd_0$, $d_0\equiv 1\pmod 4$ and $4\mid y$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then

$$(a+\sqrt{a^2+1})^{\frac{p-1}{4}}\equiv(-1)^{\frac{d}{4}+\frac{y}{4}}\pmod p.$$

$\star$ Let $p\equiv 1,9\pmod{40}$ be a prime and so $p=c^2+d^2=x^2+10y^2$ with $c,d,x,y\in\mathbb{Z}$. Suppose $c\equiv 1\pmod 4$, $d=2^rd_0$ and $d_0\equiv 1\pmod 4$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then

$$(-5)^{\frac{p-1}{8}}\equiv\begin{cases}\pm(-1)^{\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}}\pmod p & \text{if } \frac{2c-d}{x}\equiv\pm 2\pmod 5,\\ \pm(-1)^{\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}}\frac{c}{d}\pmod p & \text{if } \frac{2c-d}{x}\equiv\pm 1\pmod 5.\end{cases}$$

Throughout this paper, if $n\in\mathbb{Z}$, $2^\alpha\mid n$ and $2^{\alpha+1}\nmid n$, we then write that $2^\alpha\parallel n$.

2. Basic lemmas.

Lemma 2.1 ([S4, Proposition 2.1]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. Then*

$$\Big(\frac{i}{a+bi}\Big)_4 = i^{\frac{a^2+b^2-1}{4}} = (-1)^{\frac{a^2-1}{8}} i^{(1-(-1)^{\frac{b}{2}})/2}$$

and

$$\Big(\frac{1+i}{a+bi}\Big)_4 = \begin{cases} i^{((-1)^{\frac{a-1}{2}}(a-b)-1)/4} & \text{if } 4 \mid b, \\ i^{\frac{(-1)^{\frac{a-1}{2}}(b-a)-1}{4}-1} & \text{if } 2 \parallel b. \end{cases}$$

Lemma 2.2 ([S4, Proposition 2.2]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. Then*

$$\Big(\frac{-1}{a+bi}\Big)_4 = (-1)^{\frac{b}{2}} \quad \text{and} \quad \Big(\frac{2}{a+bi}\Big)_4 = i^{(-1)^{\frac{a-1}{2}}\frac{b}{2}}.$$

Lemma 2.3 ([S4, Proposition 2.3]). *Let $a, b, c, d \in \mathbb{Z}$ with $2 \nmid ac$, $2 \mid b$ and $2 \mid d$. If $a+bi$ and $c+di$ are relatively prime elements of $\mathbb{Z}[i]$, we have the following general law of quartic reciprocity:*

$$\Big(\frac{a+bi}{c+di}\Big)_4 = (-1)^{\frac{b}{2}\cdot\frac{c-1}{2}+\frac{d}{2}\cdot\frac{a+b-1}{2}}\Big(\frac{c+di}{a+bi}\Big)_4.$$

In particular, if $4 \mid b$, then

$$\Big(\frac{a+bi}{c+di}\Big)_4 = (-1)^{\frac{a-1}{2}\cdot\frac{d}{2}}\Big(\frac{c+di}{a+bi}\Big)_4.$$

Lemma 2.4 ([E], [S1, Lemma 2.1]). *Let $a, b, m \in \mathbb{Z}$ with $2 \nmid m$ and $(m, a^2+b^2) = 1$. Then*

$$\Big(\frac{a+bi}{m}\Big)_4^2 = \Big(\frac{a^2+b^2}{m}\Big).$$

Lemma 2.5 ([S3, Lemma 4.3]). *Let $a, b \in \mathbb{Z}$ with $2 \nmid a$ and $2 \mid b$. For any integer x with $(x, a^2+b^2) = 1$ we have*

$$\Big(\frac{x^2}{a+bi}\Big)_4 = \Big(\frac{x}{a^2+b^2}\Big).$$

For a given odd prime p let $\mathbb{Z}_p$ denote the set of those rational numbers whose denominator is not divisible by p. Following [S1,S2] we define

$$Q_r(p) = \Big\{k \;\Big|\; k \in \mathbb{Z}_p,\ \Big(\frac{k+i}{p}\Big)_4 = i^r\Big\} \quad \text{for} \quad r = 0, 1, 2, 3.$$

Lemma 2.6 ([S1, Theorem 2.3]). *Let p be an odd prime, $r\in\{0,1,2,3\}$, $k\in\mathbb{Z}_p$ and $k^2+1\not\equiv 0 \pmod p$.*

(i) *If $p\equiv 1 \pmod 4$ and $t^2\equiv -1 \pmod p$ with $t\in\mathbb{Z}_p$, then $k\in Q_r(p)$ if and only if $(\frac{k+t}{k-t})^{(p-1)/4}\equiv t^r \pmod p$.*

(ii) *If $p\equiv 3 \pmod 4$, then $k\in Q_r(p)$ if and only if $(\frac{k-i}{k+i})^{(p+1)/4}\equiv i^r \pmod p$.*

Lemma 2.7 ([S5, Lemma 2.9]). *Suppose $c,d,m,x\in\mathbb{Z}$, $2\nmid m$, $x^2\equiv c^2+d^2 \pmod m$ and $(m,x(x+d))=1$. Then*

$$\Big(\frac{c+di}{m}\Big)_4=\Big(\frac{x(x+d)}{m}\Big).$$

Lemma 2.8. *Let p be a prime of the form $8k+1$, $q\in\mathbb{Z}$, $2\nmid q$ and $p\nmid q$. Suppose that $p=c^2+d^2=x^2+2qy^2$ with $c,d,x,y\in\mathbb{Z}$, $c\equiv 1 \pmod 4$, $d=2^rd_0$ and $d_0\equiv 1 \pmod 4$. If $\big(\frac{x/y}{c+di}\big)_4=(-1)^{s+\frac{p-1}{8}}i^n$, then*

$$(-q)^{\frac{p-1}{8}}\equiv\begin{cases}(-1)^s(\frac{d}{c})^{n-1} \pmod p & \text{if } 8\mid d-4,\\ (-1)^{s+\frac{d}{8}}(\frac{d}{c})^n \pmod p & \text{if } 8\mid d.\end{cases}$$

Proof. As $c\equiv 1 \pmod 4$ and $4\mid d$ we see that $c+di$ is primary in $\mathbb{Z}[i]$. Since $i\equiv d/c \pmod{c+di}$ we have

$$\Big(\frac{x}{y}\Big)^{\frac{p-1}{4}}\equiv\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{s+\frac{p-1}{8}}i^n\equiv(-1)^{s+\frac{p-1}{8}}\Big(\frac{d}{c}\Big)^n \pmod{c+di}.$$

As $(\frac{x}{y})^2\equiv -q \pmod p$ and the norm of $c+di$ is p, from the above we deduce that

$$(-2q)^{\frac{p-1}{8}}\equiv\Big(\frac{x}{y}\Big)^{\frac{p-1}{4}}\equiv(-1)^{s+\frac{p-1}{8}}\Big(\frac{d}{c}\Big)^n \pmod p.$$

By [L] or [HW, (1.4) and (1.5)],

$$(2.1)\qquad (-2)^{\frac{p-1}{8}}\equiv\Big(\frac{c}{d}\Big)^{-\frac{d}{4}}\equiv\begin{cases}(\frac{c}{d})^{-d_0}\equiv(\frac{c}{d})^{-1}=\frac{d}{c} \pmod p & \text{if } 8\mid d-4,\\ (-1)^{\frac{d}{8}} \pmod p & \text{if } 8\mid d.\end{cases}$$

Thus the result follows.

3. Congruences for $q^{[p/8]} \pmod p$ with $4p=x^2+qy^2$.

Theorem 3.1. *Let p be a prime of the form $4k+1$, $q\in\mathbb{Z}$, $q\equiv 3 \pmod 8$ and $p\nmid q$. Suppose that $p=c^2+d^2$, $4p=x^2+qy^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv y\equiv 1 \pmod 4$, $(c,x+2d)=1$ and $(\frac{2c/(x+2d)+i}{q})_4=i^k$. Then*

$$(-q)^{[p/8]}\equiv\begin{cases}(-1)^{[\frac{x}{4}]+\frac{q-3}{8}}(\frac{d}{c})^{k-1} \pmod p & \text{if } 8\mid p-1,\\ -(-1)^{\frac{x^2-1}{8}+\frac{q-3}{8}}(\frac{d}{c})^{k-1}\frac{y}{x} \pmod p & \text{if } 8\mid p-5.\end{cases}$$

Proof. Since $4p = x^2 + qy^2$ and $y \equiv 1 \pmod 4$ we see that $2 \nmid x$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, (2c)^2 + (x + 2d)^2) = (x, 4p) = 1$. As $qy^2 = (2c)^2 + (x + 2d)(2d - x)$ we see that $(qy, x + 2d) \mid 4c^2$ and so $(qy, x + 2d) = 1$. Also,

$$\begin{aligned}
&(qy^2, (2c)^2 + (x + 2d)^2) \\
&= ((2c)^2 + (x + 2d)^2 - 2x(x + 2d), (2c)^2 + (x + 2d)^2) = (2x(x + 2d), (2c)^2 + (x + 2d)^2) \\
&= (x + 2d, (2c)^2 + (x + 2d)^2) = (x + 2d, 4c^2) = 1.
\end{aligned}$$

Now using Lemmas 2.1-2.5 and the fact that $(\frac{a}{n})_4 = 1$ for $a, n \in \mathbb{Z}$ with $2 \nmid n$ and $(a, n) = 1$ we see that

$$\begin{aligned}
i^k &= \Big(\frac{2c + (x + 2d)i}{q}\Big)_4 = \Big(\frac{i}{q}\Big)_4 \Big(\frac{x + 2d - 2ci}{q}\Big)_4 \\
&= (-1)^{\frac{q^2-1}{8}} \cdot (-1)^{\frac{q-1}{2}} \Big(\frac{q}{x + 2d - 2ci}\Big)_4 = \Big(\frac{qy^2}{x + 2d - 2ci}\Big)_4 \Big(\frac{y^2}{x + 2d - 2ci}\Big)_4 \\
&= \Big(\frac{(x + 2d)^2 + (2c)^2 - 2x(x + 2d)}{x + 2d - 2ci}\Big)_4 \Big(\frac{y}{(x + 2d)^2 + 4c^2}\Big) \\
&= \Big(\frac{-2x(x + 2d)}{x + 2d - 2ci}\Big)_4 \Big(\frac{y}{(x + 2d)^2 + 4c^2}\Big) \\
&= -\Big(\frac{2}{x + 2d - 2ci}\Big)_4 \Big(\frac{x(x + 2d)}{x + 2d - 2ci}\Big)_4 \Big(\frac{y}{(x + 2d)^2 + 4c^2}\Big) \\
&= -i^{(-1)^{(x+1)/2}} (-1)^{\frac{x(x+2d)-1}{2}} \Big(\frac{x + 2d - 2ci}{x(x + 2d)}\Big)_4 \Big(\frac{(x + 2d)^2 + 4c^2}{y}\Big) \\
&= (-1)^{\frac{x-1}{2}} i \Big(\frac{2(d - ci)}{x}\Big)_4 \Big(\frac{-2ci}{x + 2d}\Big)_4 \Big(\frac{2x(x + 2d) + qy^2}{y}\Big) \\
&= (-1)^{\frac{x-1}{2}} i \Big(\frac{d - ci}{x}\Big)_4 \Big(\frac{i}{x + 2d}\Big)_4 \Big(\frac{2x(x + 2d)}{y}\Big).
\end{aligned}$$

Thus, applying Lemma 2.7 we see that

$$\begin{aligned}
i^k &= (-1)^{\frac{x-1}{2}} i \Big(\frac{-i}{x}\Big)_4 \Big(\frac{c + di}{x}\Big)_4 (-1)^{\frac{(x+2d)^2-1}{8}} \cdot (-1)^{\frac{y^2-1}{8}} \Big(\frac{x(x + 2d)}{y}\Big) \\
&= (-1)^{\frac{x-1}{2}} i \cdot (-1)^{\frac{x^2-1}{8}} \Big(\frac{c + di}{x}\Big)_4 (-1)^{\frac{x^2-1}{8} + \frac{d}{2}} \cdot (-1)^{\frac{4p - x^2 - q}{8}} \Big(\frac{\frac{x}{2}(\frac{x}{2} + d)}{y}\Big) \\
&= (-1)^{\frac{x-1}{2} + \frac{d}{2}} i \cdot (-1)^{\frac{x-1}{2} \cdot \frac{d}{2}} \Big(\frac{x}{c + di}\Big)_4 (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} \Big(\frac{c + di}{y}\Big)_4 \\
&= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} i \Big(\frac{x}{c + di}\Big)_4 (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} \Big(\frac{y}{c + di}\Big)_4 \\
&= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} \cdot (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} i \Big(\frac{x}{c + di}\Big)_4 \Big(\frac{y^{-1}}{c + di}\Big)_4 \Big(\frac{y^2}{c + di}\Big)_4 \\
&= -(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)} \cdot (-1)^{\frac{x^2-1}{8} + \frac{q-3}{8}} i \Big(\frac{x/y}{c + di}\Big)_4 \Big(\frac{y}{c^2 + d^2}\Big).
\end{aligned}$$

As $\left(\frac{y}{c^2+d^2}\right)=\left(\frac{c^2+d^2}{y}\right)=\left(\frac{4(c^2+d^2)}{y}\right)=\left(\frac{x^2+qy^2}{y}\right)=\left(\frac{x^2}{y}\right)=1$, from the above we deduce that

$$\left(\frac{x/y}{c+di}\right)_4=-(-1)^{\frac{x+1}{2}(\frac{d}{2}+1)}\cdot(-1)^{\frac{x^2-1}{8}+\frac{q-3}{8}}i^{k-1}.$$

Clearly $(-1)^{\frac{d}{2}}=(-1)^{\frac{p-1}{4}}$ and $i\equiv d/c\pmod{c+di}$. Since $c+di$ or $-c-di$ is primary in $\mathbb{Z}[i]$, we have

$$\left(\frac{x}{y}\right)^{(p-1)/4}\equiv\left(\frac{x/y}{c+di}\right)_4\equiv-(-1)^{\frac{x+1}{2}\cdot\frac{p-5}{4}+\frac{x^2-1}{8}+\frac{q-3}{8}}\left(\frac{d}{c}\right)^{k-1}\pmod{c+di}$$

and so

$$\left(\frac{x}{y}\right)^{(p-1)/4}\equiv-(-1)^{\frac{x+1}{2}\cdot\frac{p-5}{4}+\frac{x^2-1}{8}+\frac{q-3}{8}}\left(\frac{d}{c}\right)^{k-1}\pmod{p}.$$

Note that $(x/y)^2\equiv -q\pmod{p}$. We then have

$$(-q)^{[p/8]}\equiv\begin{cases}(\frac{x}{y})^{\frac{p-1}{4}}\equiv(-1)^{\frac{x-1}{2}+\frac{x^2-1}{8}+\frac{q-3}{8}}(\frac{d}{c})^{k-1}\pmod{p} & \text{if } 8\mid p-1,\\ (\frac{x}{y})^{\frac{p-1}{4}}\frac{y}{x}\equiv-(-1)^{\frac{x^2-1}{8}+\frac{q-3}{8}}(\frac{d}{c})^{k-1}\frac{y}{x}\pmod{p} & \text{if } 8\mid p-5.\end{cases}$$

This yields the result.

Corollary 3.1. *Let p be a prime of the form $12k+1$, $p=c^2+d^2$, $4p=x^2+27y^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv y\equiv 1\pmod 4$ and $(c,x+2d)=1$.*

(i) *If $p\equiv 1\pmod{24}$, then*

$$(-3)^{\frac{p-1}{8}}\equiv\begin{cases}(-1)^{[\frac{x}{4}]}\frac{d}{c}\pmod{p} & \text{if } c\equiv 0\pmod 3,\\ \pm(-1)^{[\frac{x}{4}]}\pmod{p} & \text{if } x-d\equiv\pm c\pmod 3.\end{cases}$$

(ii) *If $p\equiv 13\pmod{24}$, then*

$$(-3)^{\frac{p-5}{8}}\equiv\begin{cases}-(-1)^{\frac{x^2-1}{8}}\frac{3dy}{cx}\pmod{p} & \text{if } c\equiv 0\pmod 3,\\ \pm(-1)^{\frac{x^2-1}{8}}\frac{3y}{x}\pmod{p} & \text{if } x-d\equiv\pm c\pmod 3.\end{cases}$$

Proof. Clearly

$$\left(\frac{i}{3}\right)_4=-1,\quad\left(\frac{1+i}{3}\right)_4=-i,\quad\left(\frac{-1+i}{3}\right)_4=i.$$

and

$$(-3)^{[\frac{p}{8}]}=3^{4[\frac{p}{8}]}(-27)^{-[\frac{p}{8}]}\equiv 3^{(-1)^{(p-1)/4}-1}(-27)^{-[\frac{p}{8}]}\pmod{p}.$$

Now taking $q=27$ in Theorem 3.1 and then applying the above we deduce the result.

Theorem 3.2. *Let p be a prime of the form $4k+1$, $q \in \mathbb{Z}$, $q \equiv 3 \pmod 8$ and $p \nmid q$. Suppose that $p = c^2 + d^2$, $4p = x^2 + qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod 4$, $(d, x+2c) = 1$ and $(\frac{-2d/(x+2c)+i}{q})_4 = i^k$. Then*

$$(-q)^{[p/8]} \equiv \begin{cases} (-1)^{[\frac{x}{4}]}(\frac{d}{c})^k \pmod p & \text{if } 8 \mid p-1, \\ -(-1)^{\frac{x^2-1}{8}}(\frac{d}{c})^k \frac{y}{x} \pmod p & \text{if } 8 \mid p-5. \end{cases}$$

Proof. Since $4p = x^2 + qy^2$ and $y \equiv 1 \pmod 4$ we see that $2 \nmid x$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, (2d)^2 + (x+2c)^2) = (x, 4p) = 1$. As $qy^2 = (2d)^2 + (x+2c)(2c-x)$ we see that $(qy, x+2c) \mid (2d)^2$ and so $(qy, x+2c) = 1$. Also,

$$\begin{aligned} &(qy^2, (2d)^2 + (x+2c)^2) \\ &= ((2d)^2 + (x+2c)^2 - 2x(x+2c), (2d)^2 + (x+2c)^2) = (2x(x+2c), (2d)^2 + (x+2c)^2) \\ &= (x+2c, (2d)^2 + (x+2c)^2) = (x+2c, 4d^2) = 1. \end{aligned}$$

Now using Lemmas 2.1-2.5 and the fact that $(\frac{a}{n})_4 = 1$ for $a, n \in \mathbb{Z}$ with $2 \nmid n$ and $(a, n) = 1$ we see that

$$\begin{aligned} i^k &= \Big(\frac{-2d + (x+2c)i}{q}\Big)_4 = \Big(\frac{i}{q}\Big)_4 \Big(\frac{x+2c+2di}{q}\Big)_4 \\ &= (-1)^{\frac{q^2-1}{8}} \Big(\frac{q}{x+2c+2di}\Big)_4 = -\Big(\frac{qy^2}{x+2c+2di}\Big)_4 \Big(\frac{y^2}{x+2c+2di}\Big)_4 \\ &= -\Big(\frac{(x+2c)^2 + (2d)^2 - 2x(x+2c)}{x+2c+2di}\Big)_4 \Big(\frac{y}{(x+2c)^2+4d^2}\Big) \\ &= -\Big(\frac{2}{x+2c+2di}\Big)_4 \Big(\frac{x(x+2c)}{x+2c+2di}\Big)_4 \Big(\frac{y}{(x+2c)^2+4d^2}\Big) \\ &= -(-1)^{\frac{d}{2}} \Big(\frac{x+2c+2di}{x(x+2c)}\Big)_4 \Big(\frac{(x+2c)^2+4d^2}{y}\Big) \\ &= -(-1)^{\frac{p-1}{4}} \Big(\frac{2(c+di)}{x}\Big)_4 \Big(\frac{2di}{x+2c}\Big)_4 \Big(\frac{2x(x+2c)+qy^2}{y}\Big) \\ &= -(-1)^{\frac{p-1}{4}} \Big(\frac{c+di}{x}\Big)_4 \Big(\frac{i}{x+2c}\Big)_4 \Big(\frac{2x(x+2c)}{y}\Big). \end{aligned}$$

Thus, applying Lemma 2.7 we see that

$$\begin{aligned}
i^k &= -(-1)^{\frac{p-1}{4}}\Big(\frac{c+di}{x}\Big)_4(-1)^{\frac{(x+2c)^2-1}{8}}\Big(\frac{2}{y}\Big)\Big(\frac{x(x+2c)}{y}\Big)\\
&= -(-1)^{\frac{p-1}{4}}\Big(\frac{c+di}{x}\Big)_4(-1)^{\frac{x^2-1}{8}+\frac{x+c}{2}}\Big(\frac{i}{y}\Big)_4\Big(\frac{\frac{x}{2}(\frac{x}{2}+c)}{y}\Big)\\
&= -(-1)^{\frac{p-1}{4}}\cdot(-1)^{\frac{x-1}{2}\cdot\frac{d}{2}}\Big(\frac{x}{c+di}\Big)_4(-1)^{\frac{x^2-1}{8}+\frac{x+1}{2}}\Big(\frac{i}{y}\Big)_4\Big(\frac{d+ci}{y}\Big)_4\\
&= -(-1)^{(1+\frac{x-1}{2})(1+\frac{p-1}{4})}\cdot(-1)^{\frac{x^2-1}{8}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{-c+di}{y}\Big)_4\\
&= -(-1)^{\frac{x+1}{2}\cdot\frac{p+3}{4}+\frac{x^2-1}{8}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{c+di}{y}\Big)_4^{-1}\\
&= -(-1)^{\frac{x+1}{2}\cdot\frac{p+3}{4}+\frac{x^2-1}{8}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{y}{c+di}\Big)_4^{-1}\\
&= -(-1)^{\frac{x+1}{2}\cdot\frac{p+3}{4}+\frac{x^2-1}{8}}\Big(\frac{x/y}{c+di}\Big)_4.
\end{aligned}$$

Clearly $(-1)^{\frac{d}{2}} = (-1)^{\frac{p-1}{4}}$ and $i \equiv d/c \pmod{c+di}$. Since $c+di$ or $-c-di$ is primary in $\mathbb{Z}[i]$, we have

$$\Big(\frac{x}{y}\Big)^{(p-1)/4} \equiv \Big(\frac{x/y}{c+di}\Big)_4 \equiv -(-1)^{\frac{x+1}{2}\cdot\frac{p+3}{4}+\frac{x^2-1}{8}}\Big(\frac{d}{c}\Big)^k \pmod{c+di}$$

and so

$$\Big(\frac{x}{y}\Big)^{(p-1)/4} \equiv -(-1)^{\frac{x+1}{2}\cdot\frac{p+3}{4}+\frac{x^2-1}{8}}\Big(\frac{d}{c}\Big)^k \pmod{p}.$$

Note that $(x/y)^2 \equiv -q \pmod{p}$. We then have

$$(-q)^{[p/8]} \equiv \begin{cases} (\frac{x}{y})^{\frac{p-1}{4}} \equiv -(-1)^{\frac{x+1}{2}+\frac{x^2-1}{8}}(\frac{d}{c})^k = (-1)^{[\frac{x}{4}]}(\frac{d}{c})^k \pmod{p} & \text{if } 8 \mid p-1,\\ (\frac{x}{y})^{\frac{p-1}{4}}\frac{y}{x} \equiv -(-1)^{\frac{x^2-1}{8}}(\frac{d}{c})^k\frac{y}{x} \pmod{p} & \text{if } 8 \mid p-5. \end{cases}$$

This is the result.

Remark 3.1 We note that the k in Theorem 3.1 depends only on $\frac{c}{x+2d} \pmod{q}$, and the k in Theorem 3.2 depends only on $\frac{d}{x+2c} \pmod{q}$.

Theorem 3.3. *Let p and q be primes such that $p \equiv 1 \pmod 4$ and $q \equiv 3 \pmod 8$. Suppose $p = c^2+d^2$, $4p = x^2+qy^2$, $c,d,x,y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod 4$, and $(\frac{2(c-di)}{x})^{\frac{q+1}{4}} \equiv i^m \pmod q$. Assume $(c, x+2d) = 1$ or $(d, x+2c) = 1$. Then*

$$(-q)^{[p/8]} \equiv \begin{cases} (-1)^{[\frac{x}{4}]+1}(\frac{d}{c})^m \pmod{p} & \text{if } p \equiv 1 \pmod 8,\\ (-1)^{\frac{x^2-1}{8}}(\frac{d}{c})^m\frac{y}{x} \pmod{p} & \text{if } p \equiv 5 \pmod 8. \end{cases}$$

Proof. Clearly $q \nmid x$ and x is odd. We first assume $(c, x+2d) = 1$. By the proof of Theorem 3.1, $(q, (x+2d)(c^2+(x+2d)^2)) = 1$. It is easily seen that $\frac{2c/(x+2d)-i}{2c/(x+2d)+i} = \frac{2c-(x+2d)i}{2c+(x+2d)i} \equiv \frac{2(c-di)}{ix} \pmod q$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned}
&\Big(\frac{2c/(x+2d)+i}{q}\Big)_4 = i^k \\
&\iff \frac{2c}{x+2d} \in Q_k(q) \iff \Big(\frac{\frac{2c}{x+2d}-i}{\frac{2c}{x+2d}+i}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \\
&\iff \Big(\frac{2(c-di)}{ix}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \iff \Big(\frac{2(c-di)}{x}\Big)^{\frac{q+1}{4}} \equiv i^{\frac{q+1}{4}+k} \pmod q.
\end{aligned}$$

Since $(\frac{2(c-di)}{x})^{\frac{q+1}{4}} \equiv i^m \pmod q$, from the above we deduce that

$$\Big(\frac{2c/(x+2d)+i}{q}\Big)_4 = i^{m-\frac{q+1}{4}} = (-1)^{\frac{q-3}{8}} i^{m-1}.$$

Now, applying Theorem 3.1 we derive the result.

Now we assume $(d, x+2c) = 1$. By the proof of Theorem 3.2, $(q, x+2c) = (q, d^2+(x+2c)^2) = 1$. It is easily seen that $\frac{2d+(x+2c)i}{2d-(x+2c)i} \equiv \frac{2(c-di)}{-x} \pmod q$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned}
&\Big(\frac{-2d/(x+2c)+i}{q}\Big)_4 = i^k \\
&\iff -\frac{2d}{x+2c} \in Q_k(q) \iff \Big(\frac{-\frac{2d}{x+2c}-i}{-\frac{2d}{x+2c}+i}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \\
&\iff \Big(\frac{2d+(x+2c)i}{2d-(x+2c)i}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \iff \Big(\frac{2(c-di)}{-x}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \\
&\iff \Big(\frac{2(c-di)}{x}\Big)^{\frac{q+1}{4}} \equiv i^{\frac{q+1}{2}+k} \pmod q.
\end{aligned}$$

Since $(\frac{2(c-di)}{x})^{\frac{q+1}{4}} \equiv i^m \pmod q$, from the above we deduce that $(\frac{-2d/(x+2c)+i}{q})_4 = i^{m-\frac{q+1}{2}} = (-1)^{\frac{q+1}{4}} i^m = -i^m$. Now applying Theorem 3.2 we deduce the result. The proof is now complete.

Theorem 3.4. *Let p and q be primes such that $p \equiv 1 \pmod 4$ and $q \equiv 3 \pmod 8$. Suppose $p = c^2+d^2$, $4p = x^2+qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv y \equiv 1 \pmod 4$, and $q \mid cd$. Assume that $(c, x+2d) = 1$ or $(d, x+2c) = 1$. Then*

$$(-q)^{[\frac{p}{8}]} \equiv \begin{cases}
\pm(-1)^{[\frac{x}{4}]+1} \pmod p & \text{if } p \equiv 1 \pmod 8 \text{ and } x \equiv \pm 2c \pmod q, \\
\pm(-1)^{\frac{q-3}{8}+[\frac{x}{4}]}\frac{d}{c} \pmod p & \text{if } p \equiv 1 \pmod 8 \text{ and } x \equiv \pm 2d \pmod q, \\
\pm(-1)^{\frac{x^2-1}{8}}\frac{y}{x} \pmod p & \text{if } p \equiv 5 \pmod 8 \text{ and } x \equiv \pm 2c \pmod q, \\
\mp(-1)^{\frac{q-3}{8}+\frac{x^2-1}{8}}\frac{dy}{cx} \pmod p & \text{if } p \equiv 5 \pmod 8 \text{ and } x \equiv \pm 2d \pmod q,
\end{cases}$$

Proof. As $(\frac{x}{2})^2 \equiv c^2+d^2 \pmod q$ we see that $x \equiv \pm 2c \pmod q \Leftrightarrow q \mid d$ and that $x \equiv \pm 2d \pmod q \Leftrightarrow q \mid c$. Thus,

$$\Big(\frac{c-di}{x/2}\Big)_4^{\frac{q+1}{4}} \equiv \begin{cases} (\pm 1)^{\frac{q+1}{4}} = \pm 1 \pmod q & \text{if } x \equiv \pm 2c \pmod q, \\ (\mp i)^{\frac{q-1}{4}} = \mp(-1)^{\frac{q-3}{8}} i \pmod q & \text{if } x \equiv \pm 2d \pmod q. \end{cases}$$

Now applying Theorem 3.3 we derive the result.

Theorem 3.5. *Let p be a prime of the form $4k+1$ and $(\frac{-11}{p}) = 1$, $p = c^2+d^2$, $4p = x^2+11y^2$, $c,d,x,y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0 (2 \nmid d_0)$, $y = 2^t y_0$ and $y_0 \equiv 1 \pmod 4$. Assume that $(c, x+2d) = 1$ or $(d_0, x+2c) = 1$.*

(i) *If $p \equiv 1 \pmod 8$, then*

$$(-11)^{[\frac{p}{8}]} \equiv \begin{cases} \pm(-1)^{[\frac{x}{4}]} \pmod p & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \pm(-1)^{[\frac{x}{4}]} \frac{d}{c} \pmod p & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}, \\ \mp(-1)^{[\frac{x}{8}]+\frac{y}{8}} \pmod p & \text{if } 2 \mid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{[\frac{x}{8}]+\frac{y}{8}} \frac{d}{c} \pmod p & \text{if } 2 \mid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

(ii) *If $p \equiv 5 \pmod 8$, then*

$$(-11)^{[\frac{p}{8}]} \equiv \begin{cases} \mp(-1)^{\frac{x^2-1}{8}} \frac{y}{x} \pmod p & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\frac{x^2-1}{8}} \frac{dy}{cx} \pmod p & \text{if } 2 \nmid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}, \\ \mp(-1)^{\frac{p-5}{8}} \frac{y}{x} \pmod p & \text{if } 2 \mid x \text{ and } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp(-1)^{\frac{p-5}{8}} \frac{dy}{cx} \pmod p & \text{if } 2 \mid x \text{ and } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

Proof. As

$$\Big(\frac{x}{2}\Big)^2 \equiv c^2+d^2 \pmod{11} \quad \text{and} \quad (c-di)^3 = c(c^2-3d^2) + d(d^2-3c^2)i,$$

we see that

$$\Big(\frac{c-di}{x/2}\Big)_4^3 \equiv \begin{cases} \mp 1 \pmod p & \text{if } x \equiv \pm 4c, \pm 9c \pmod{11}, \\ \mp i \pmod p & \text{if } x \equiv \pm 4d, \pm 9d \pmod{11}. \end{cases}$$

When $2 \nmid x$, from the above and Theorem 3.3 (with $q = 11$) we deduce the result. When $2 \mid x$ and $p \equiv 1 \pmod 8$, we have $8 \mid y$ and so $(-1)^{\frac{p-1}{8}+\frac{x/2-1}{2}} = (-1)^{\frac{(x/2)^2-1}{8}+\frac{x/2-1}{2}} = (-1)^{[\frac{x}{8}]}$. Thus, applying the above and [S5, Theorem 4.1 (with $q = 11$)] we obtain the result.

Remark 3.2 Let p be a prime of the form $4k+1$, $q \in \mathbb{Z}$, $q \equiv 3 \pmod 8$, $p \nmid q$, $p = c^2+d^2$, $4p = x^2+qy^2$, $c,d,x,y \in \mathbb{Z}$, $2 \nmid xy$, $c \equiv 1 \pmod 4$, we conjecture that one can always choose the sign of x such that $(c, x+2d) = 1$ or $(d, x+2c) = 1$. Thus the condition $(c, x+2d) = 1$ or $(d, x+2c) = 1$ in Theorems 3.1-3.5 can be canceled. See also [S5, Remark 4.1].

4. Congruences for $(-q)^{(p-1)/8} \pmod p$ with $p = x^2 + 2qy^2$.

Theorem 4.1. *Let p be a prime of the form $8k+1$, $q \in \mathbb{Z}$, $2 \nmid q$ and $p \nmid q$. Suppose that $p = c^2 + d^2 = x^2 + 2qy^2$ with $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$, $d_0 \equiv 1 \pmod 4$ and $(c, x+d) = 1$. Assume that $(\frac{c/(x+d)+i}{q})_4 = i^k$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8}+\frac{d}{4}+\frac{y}{2}}(\frac{d}{c})^k \pmod p & \text{if } q \equiv 1 \pmod 8, \\ (-1)^{\frac{q-3}{8}+\frac{x-1}{2}}(\frac{d}{c})^{k+1} \pmod p & \text{if } q \equiv 3 \pmod 8, \\ (-1)^{\frac{q-5}{8}+\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}}(\frac{d}{c})^{k-1} \pmod p & \text{if } q \equiv 5 \pmod 8, \\ (-1)^{\frac{q+1}{8}}(\frac{d}{c})^k \pmod p & \text{if } q \equiv 7 \pmod 8. \end{cases}$$

Proof. Suppose $y = 2^t y_0$ ($2 \nmid y_0$). We may assume $y_0 \equiv 1 \pmod 4$. Since $p = c^2 + d^2 = x^2 + 2qy^2 \equiv 1 \pmod 8$ we see that $2 \nmid x$, $2 \mid y$, $4 \mid d$, $(x, y) = 1$ and $p \nmid x$. Thus $(x, c^2 + (x+d)^2) = (x, p) = 1$. As $2qy^2 = c^2 + (d+x)(d-x)$ we see that $(qy, x+d) \mid c^2$ and so $(qy, x+d) = 1$. Also,

$$\begin{aligned} &(qy^2, (c^2 + (x+d)^2)/2) \\ &= ((c^2 + (x+d)^2)/2 - x(x+d), (c^2 + (x+d)^2)/2) = (x(x+d), (c^2 + (x+d)^2)/2) \\ &= (x(x+d), c^2 + (x+d)^2) = (x+d, c^2) = 1. \end{aligned}$$

It is easily seen that

$$c + (x+d)i = i^{\frac{1\mp 1}{2}}(1+i)\Big(\frac{x+d \pm c}{2} + \frac{\pm(x+d) - c}{2}i\Big)$$

and so

$$\Big(\frac{x+d \pm c}{2}\Big)^2 + \Big(\frac{\pm(x+d) - c}{2}\Big)^2 = \frac{c^2 + (x+d)^2}{2}.$$

Set $\varepsilon = (-1)^{\frac{x-1}{2}}$. As $4 \mid d$ and $4 \mid c-1$ we have $x+d \equiv \varepsilon \pmod 4$ and $4 \mid (\varepsilon(x+d) - c)$. Using Lemmas 2.1-2.5 and [S5, Lemma 2.10(ii)] and the above we see that

$$\begin{aligned} i^k &= \Big(\frac{c + (x+d)i}{q}\Big)_4 = \Big(\frac{i}{q}\Big)_4^{\frac{1-\varepsilon}{2}}\Big(\frac{1+i}{q}\Big)_4\Big(\frac{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i}{q}\Big)_4 \\ &= (-1)^{\frac{q-(\frac{-1}{q})}{4}\cdot\frac{x-1}{2}} i^{\frac{(\frac{-1}{q})q-1}{4}} (-1)^{\frac{q-1}{2}\cdot\frac{\varepsilon(x+d)-c}{4}} \Big(\frac{q}{\frac{x+d+\varepsilon c}{2} + \frac{\varepsilon(x+d)-c}{2}i}\Big)_4 \end{aligned}$$

and

$$\begin{aligned}
&\Big(\frac{q}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4\\
&=\Big(\frac{qy^2}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4\Big(\frac{y^2}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4\\
&=\Big(\frac{(c^2+(x+d)^2)/2-x(x+d)}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4\Big(\frac{y}{(\frac{x+d+\varepsilon c}{2})^2+(\frac{\varepsilon(x+d)-c}{2})^2}\Big)\\
&=\Big(\frac{-x(x+d)}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4\Big(\frac{y}{(c^2+(x+d)^2)/2}\Big)\\
&=(-1)^{\frac{\varepsilon(x+d)-c}{4}}\Big(\frac{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}{x(x+d)}\Big)_4(-1)^{\frac{c^2-(x+d)^2}{8}t+\frac{d}{4}t}\Big(\frac{y^{-1}}{c+di}\Big)_4.
\end{aligned}$$

Clearly,

$$(-1)^{\frac{c^2-(x+d)^2}{8}}=(-1)^{\frac{c^2-x^2-2dx}{8}}=(-1)^{\frac{c-\varepsilon x}{4}\cdot\frac{c+\varepsilon x}{2}-\frac{d}{4}\varepsilon}=(-1)^{\frac{c-\varepsilon x}{4}-\frac{d}{4}\varepsilon}=(-1)^{\frac{\varepsilon(x+d)-c}{4}}.$$

It is easily seen that

$$\begin{aligned}
&\Big(\frac{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}{x(x+d)}\Big)_4\\
&=\Big(\frac{x+d+\varepsilon c+(\varepsilon(x+d)-c)i}{x}\Big)_4\Big(\frac{x+d+\varepsilon c+(\varepsilon(x+d)-c)i}{x+d}\Big)_4\\
&=\Big(\frac{d+\varepsilon c+(\varepsilon d-c)i}{x}\Big)_4\Big(\frac{\varepsilon c-ci}{x+d}\Big)_4=\Big(\frac{(\varepsilon-i)(c+di)}{x}\Big)_4\Big(\frac{\varepsilon-i}{x+d}\Big)_4\\
&=\Big(\frac{\varepsilon-i}{x(x+d)}\Big)_4\Big(\frac{c+di}{x}\Big)_4=\Big(\frac{i^{\frac{5+\varepsilon}{2}}(1+i)}{x(x+d)}\Big)_4\Big(\frac{c+di}{x}\Big)_4\\
&=\Big(\frac{i}{x(x+d)}\Big)_4^{\frac{5+\varepsilon}{2}}\Big(\frac{1+i}{x(x+d)}\Big)_4\Big(\frac{x}{c+di}\Big)_4\\
&=(-1)^{\frac{x(x+d)-1}{4}\cdot\frac{5+\varepsilon}{2}}i^{\frac{x(x+d)-1}{4}}\Big(\frac{x}{c+di}\Big)_4\\
&=(-1)^{\frac{d}{4}\cdot\frac{x+1}{2}+\frac{x^2-1}{8}}i^{\frac{dx}{4}}\Big(\frac{x}{c+di}\Big)_4.
\end{aligned}$$

Hence

$$\begin{aligned}
\Big(\frac{q}{\frac{x+d+\varepsilon c}{2}+\frac{\varepsilon(x+d)-c}{2}i}\Big)_4&=(-1)^{\frac{\varepsilon(x+d)-c}{4}(1+t)+\frac{d}{4}t}\cdot(-1)^{\frac{d}{4}\cdot\frac{x+1}{2}+\frac{x^2-1}{8}}i^{\frac{dx}{4}}\Big(\frac{x/y}{c+di}\Big)_4\\
&=(-1)^{\frac{\varepsilon x-c}{4}(1+t)+\frac{d}{4}\cdot\frac{x-1}{2}+\frac{x^2-1}{8}}i^{\frac{dx}{4}}\Big(\frac{x/y}{c+di}\Big)_4.
\end{aligned}$$

Therefore

$$i^k=(-1)^{\frac{q-(\frac{-1}{q})}{4}\cdot\frac{x-1}{2}+\frac{q-1}{2}(\frac{\varepsilon x-c}{4}+\frac{d}{4})}i^{\frac{(\frac{-1}{q})q-1}{4}}$$
$$\times(-1)^{\frac{\varepsilon x-c}{4}(1+t)+\frac{d}{4}\cdot\frac{x-1}{2}+\frac{x^2-1}{8}}i^{\frac{dx}{4}}\Big(\frac{x/y}{c+di}\Big)_4.$$

It is clear that

$$i^{\frac{dx}{4}}=i^{\frac{d}{4}(x-1)+\frac{d}{4}}=(-1)^{\frac{d}{4}\cdot\frac{x-1}{2}}i^{\frac{d}{4}},\ (-1)^{\frac{x^2-1}{8}}=(-1)^{\frac{p-1-2qy^2}{8}}=(-1)^{\frac{p-1}{8}+\frac{y}{2}},$$
$$(-1)^{\frac{\varepsilon x-c}{4}}=(-1)^{\frac{\varepsilon x-c}{4}\cdot\frac{\varepsilon x+c}{2}}=(-1)^{\frac{x^2-c^2}{8}}=(-1)^{\frac{d^2-8q(\frac{y}{2})^2}{8}}=(-1)^{\frac{y}{2}}$$

and so $(-1)^{\frac{\varepsilon x-c}{4}(1+t)}=(-1)^{\frac{y}{2}(1+t)}=1$. Thus,

$$\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{q-(\frac{-1}{q})}{4}\cdot\frac{x-1}{2}+\frac{q-1}{2}(\frac{y}{2}+\frac{d}{4})+\frac{p-1}{8}+\frac{y}{2}}i^{k-\frac{d}{4}-\frac{q(\frac{-1}{q})-1}{4}}.$$

Now applying Lemma 2.8 we deduce the result.

Theorem 4.2. *Let p be a prime of the form $4k+1$, $q\in\mathbb{Z}$, $2\nmid q$ and $p\nmid q$. Suppose that $p=c^2+d^2=x^2+2qy^2$ with $c,d,x,y\in\mathbb{Z}$, $c\equiv 1\pmod 4$, $d=2^rd_0$, $d_0\equiv 1\pmod 4$, $(d_0,x+c)=1$ and $(\frac{-d/(x+c)+i}{q})_4=i^k$. Then*

$$(-q)^{\frac{p-1}{8}}\equiv\begin{cases}(-1)^{\frac{x+1}{2}\cdot\frac{q-1}{4}+\frac{d}{4}+\frac{y}{2}}(\frac{d}{c})^k\pmod p & \text{if } q\equiv 1\pmod 4,\\ (-1)^{\frac{x+1}{2}\cdot\frac{q+1}{4}}(\frac{d}{c})^k\pmod p & \text{if } q\equiv 3\pmod 4.\end{cases}$$

Proof. Suppose $2^m\parallel(x+c)$ and $y=2^ty_0$ $(2\nmid y_0)$. We may assume $y_0\equiv 1\pmod 4$. As $(x+c,d_0)=1$, by Lemma 2.11 we have $(qy_0,x+c)=1$ and $(qy_0^2,(x+c)^2+d^2)=1$. Note that $(x,y)^2\mid p$. We also have $(x,y)=1$. We prove the theorem by considering the following three cases:

Case 1. $m<r$. Using Lemmas 2.1-2.5 and the fact $\big(\frac{a}{q}\big)_4=1$ for $a\in\mathbb{Z}$ with $(a,q)=1$ we see that

$$\Big(\frac{d-(x+c)i}{q}\Big)_4=\Big(\frac{-2^m}{q}\Big)_4\Big(\frac{i}{q}\Big)_4\Big(\frac{\frac{x+c}{2^m}+\frac{d}{2^m}i}{q}\Big)_4=(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{q}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4$$
$$=(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{qy^2}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4\Big(\frac{y^2}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4$$
$$=(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{((x+c)^2+d^2)/2-x(x+c)}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4\Big(\frac{y}{\frac{(x+c)^2+d^2}{2^{2m}}}\Big)$$
$$=(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{-2^mx(x+c)/2^m}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4\Big(\frac{y}{\frac{(x+c)^2+d^2}{2^{2m}}}\Big).$$

By [S5, p.15],

$$\Big(\frac{y}{((x+c)^2+d^2)/2^{2m}}\Big)=(-1)^{\frac{d}{2^{m+1}}t+\frac{d}{4}t}\Big(\frac{y^{-1}}{c+di}\Big)_4.$$

We also have

$$\Big(\frac{2^m}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4=\Big(\frac{2}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4^m=i^{\frac{x+c}{2^m}\cdot\frac{dm}{2^{m+1}}}$$

and

$$\begin{aligned}\Big(\frac{-x(x+c)/2^m}{\frac{x+c}{2^m}+\frac{d}{2^m}i}\Big)_4&=(-1)^{\frac{-x(x+c)/2^m-1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{\frac{x+c}{2^m}+\frac{d}{2^m}i}{-x(x+c)/2^m}\Big)_4\\&=(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{\frac{x+c}{2^m}+\frac{d}{2^m}i}{-x}\Big)_4\Big(\frac{\frac{x+c}{2^m}+\frac{d}{2^m}i}{(x+c)/2^m}\Big)_4\\&=(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{x+c+di}{x}\Big)_4\Big(\frac{\frac{d}{2^m}i}{(x+c)/2^m}\Big)_4\\&=(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{c+di}{x}\Big)_4\Big(\frac{i}{(x+c)/2^m}\Big)_4\\&=(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}}\Big(\frac{x}{c+di}\Big)_4(-1)^{\frac{(\frac{x+c}{2^m})^2-1}{8}}.\end{aligned}$$

Thus,

$$\begin{aligned}i^k=\Big(\frac{d-(x+c)i}{q}\Big)_4&=(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}}i^{\frac{x+c}{2^m}\cdot\frac{dm}{2^{m+1}}}\\&\times(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}+\frac{(\frac{x+c}{2^m})^2-1}{8}}(-1)^{\frac{d}{2^{m+1}}t+\frac{d}{4}t}\Big(\frac{x/y}{c+di}\Big)_4.\end{aligned}\tag{4.1}$$

As $2qy^2=d^2-(x+c)^2+2c(x+c)$ we have

$$q\frac{y^2}{2^m}=2^{2r-m-1}d_0^2-2^{m-1}\Big(\frac{x+c}{2^m}\Big)^2+c\cdot\frac{x+c}{2^m}.\tag{4.2}$$

Suppose $x\equiv 1\pmod 4$. Then $m=1<r$. From (4.2) we see that $2^{2t-1}q\equiv 2^{2r-2}-1+c\cdot\frac{x+c}{2}\pmod 8$ and so $\frac{x+c}{2}\equiv c(2^{2t-1}q-2^{2r-2}+1)\pmod 8$. If $8\nmid d$, then $r=2$ and $\frac{x+c}{2}\equiv c(2^{2t-1}q-3)\pmod 8$. Thus,

$$\begin{aligned}(-1)^{\frac{(\frac{x+c}{2})^2-1}{8}}&=(-1)^{\frac{c^2-1}{8}+\frac{(2^{2t-1}q-3)^2-1}{8}}=(-1)^{\frac{c^2-1}{8}+\frac{(2^{2t-2}q-2)(2^{2t-2}q-1)}{2}}\\&=(-1)^{\frac{c^2-1}{8}+1+\frac{q+1}{2}\cdot\frac{y}{2}}=(-1)^{\frac{p-1}{8}+1+\frac{q+1}{2}\cdot\frac{y}{2}}\end{aligned}$$

and

$$(-1)^{\frac{(x+c)/2+1}{2}}i^{\frac{x+c}{2}}=(-1)^{2^{2t-2}q-1}i^{2^{2t-1}q-3}=-(-1)^{2^{2t-2}}\cdot(-1)^{2^{2t-2}}i=-i.$$

Hence, from (4.1) we deduce that

$$\begin{aligned} i^k &= (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}} i^{\frac{x+c}{2}} (-1)^{\frac{(x+c)/2+1}{2}} \cdot (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} \Big(\frac{x/y}{c+di}\Big)_4 \\ &= (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}} (-i)(-1)^{\frac{p-1}{8}+1+\frac{q+1}{2}\cdot\frac{y}{2}} \Big(\frac{x/y}{c+di}\Big)_4. \end{aligned}$$

That is,

$$\Big(\frac{x/y}{c+di}\Big)_4 = (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}+\frac{p-1}{8}+\frac{q+1}{2}\cdot\frac{y}{2}} i^{k-1}.$$

Now applying Lemma 2.8 we obtain the result.

If $8 \mid d$, from (4.2) we see that $2^{2t-1}q \equiv 2q(\frac{y}{2})^2 \equiv -1 + c\cdot\frac{x+c}{2} \pmod 8$ and so $\frac{x+c}{2} \equiv c(2^{2t-1}q+1) \pmod 8$. Thus,

$$\begin{aligned} (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} &= (-1)^{\frac{c^2-1}{8}+\frac{(2^{2t-1}q+1)^2-1}{8}} = (-1)^{\frac{c^2+d^2-1}{8}+\frac{2^{2t-2}q(2^{2t-2}q+1)}{2}} \\ &= (-1)^{\frac{p-1}{8}+\frac{q+1}{2}\cdot\frac{y}{2}}. \end{aligned}$$

Now, from (4.1) and the above we derive that

$$\begin{aligned} i^k &= (-1)^{\frac{q^2-1}{8}} i^{\frac{x+c}{2}\cdot\frac{d}{4}} (-1)^{\frac{(\frac{x+c}{2})^2-1}{8}} \Big(\frac{x/y}{c+di}\Big)_4 \\ &= (-1)^{\frac{q^2-1}{8}+\frac{d}{8}+\frac{p-1}{8}+\frac{q+1}{2}\cdot\frac{y}{2}} \Big(\frac{x/y}{c+di}\Big)_4. \end{aligned}$$

Now applying Lemma 2.8 we obtain

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{q^2-1}{8}+\frac{q+1}{2}\cdot\frac{y}{2}} \Big(\frac{d}{c}\Big)^k = \begin{cases} (-1)^{\frac{q-1}{4}+\frac{d}{4}+\frac{y}{2}} (\frac{d}{c})^k \pmod p & \text{if } 4 \mid q-1, \\ (\frac{d}{c})^k \pmod p & \text{if } 4 \mid q-3. \end{cases}$$

This is the result.

Now assume $r > m \ge 2$. Then $x \equiv 3 \pmod 4$, $2r-m-1 \ge 2(m+1)-m-1 = m+1 \ge 3$ and so $q\frac{y^2}{2^m} \equiv -2^{m-1} + c\cdot\frac{x+c}{2^m} \pmod 8$. Hence $2^m \parallel y^2$, $m = 2t$ and so $q \equiv -2^{m-1} + c\cdot\frac{x+c}{2^m} \pmod 8$. That is, $\frac{x+c}{2^m} \equiv c(2^{m-1}+q) \pmod 8$. Thus,

$$\begin{aligned} &(-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\frac{d}{2^{m+1}}} (-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{d}{2^{m+1}}+\frac{(\frac{x+c}{2^m})^2-1}{8}} \cdot (-1)^{\frac{d}{2^{m+1}}t+\frac{d}{4}t} i^{\frac{x+c}{2^m}\cdot\frac{dm}{2^{m+1}}} \\ &= (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot\frac{d}{2^{m+1}}} (-1)^{\frac{-(2^{m-1}+q)+1}{2}\cdot 2^{r-m-1}d_0+\frac{c^2(2^{m-1}+q)^2-1}{8}} (-1)^{\frac{d}{2^{m+1}}t+\frac{d}{4}t} (-1)^{\frac{dt}{2^{m+1}}} \\ &= (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot 2^{r-m-1}d_0} \cdot (-1)^{(2^{m-2}+\frac{q-1}{2})2^{r-m-1}+\frac{c^2-1}{8}+\frac{(2^{m-1}+q)^2-1}{8}} \cdot (-1)^{\frac{d}{4}t} \\ &= (-1)^{\frac{q^2-1}{8}+\frac{q-1}{2}\cdot 2^{r-m-1}} \cdot (-1)^{(2^{m-2}+\frac{q-1}{2})2^{r-m-1}+\frac{p-1}{8}+\frac{q^2-1}{8}+2^{m-3}(2^{m-2}+q)} \\ &= (-1)^{2^{r-3}+\frac{p-1}{8}+2^{m-3}(2^{m-2}+q)} = \begin{cases} (-1)^{\frac{d}{8}+\frac{p-1}{8}+\frac{q+1}{2}} & \text{if } m = 2, \\ (-1)^{\frac{p-1}{8}} & \text{if } m > 2. \end{cases} \end{aligned}$$

Hence, from (4.1) and the above we get

$$\Big(\frac{x/y}{c+di}\Big)_4=\begin{cases}(-1)^{\frac{d}{8}+\frac{p-1}{8}+\frac{q+1}{2}}i^k & \text{if } r>m=2,\\ (-1)^{\frac{p-1}{8}}i^k & \text{if } r>m>2.\end{cases}$$

Now applying Lemma 2.8 and the fact $m=2t$ we obtain

$$\begin{aligned}(-q)^{\frac{p-1}{8}}&\equiv\begin{cases}(-1)^{\frac{q+1}{2}}(\frac{d}{c})^k \pmod p & \text{if } r>m=2,\\ (-1)^{\frac{d}{8}}(\frac{d}{c})^k=(\frac{d}{c})^k \pmod p & \text{if } r>m>2\end{cases}\\ &=(-1)^{\frac{q+1}{2}\cdot\frac{y}{2}}\Big(\frac{d}{c}\Big)^k=(-1)^{\frac{q+1}{2}(\frac{d}{4}+\frac{y}{2})}\Big(\frac{d}{c}\Big)^k \pmod p \quad \text{for} \quad r>m\ge 2.\end{aligned}$$

This yields the result.

Case 2. $m=r$. As $p\equiv 1 \pmod 8$ we have $4\mid d$, $m=r\ge 2$, $4\mid x+c$ and so $x\equiv 3 \pmod 4$. It is clear that

$$\begin{aligned}i^k&=\Big(\frac{-d/(x+c)+i}{q}\Big)_4=\Big(\frac{-\frac{d}{2^r}+\frac{x+c}{2^r}i}{q}\Big)_4\\ &=\Big(\frac{1+i}{q}\Big)_4\Big(\frac{\frac{x+c-d}{2^{r+1}}+\frac{x+c+d}{2^{r+1}}i}{q}\Big)_4=\Big(\frac{1+i}{q}\Big)_4\Big(\frac{\frac{x+c-d}{2^{r+1}}-\frac{x+c+d}{2^{r+1}}i}{q}\Big)_4^{-1}\\ &=\Big(\frac{i}{q}\Big)_4\Big(\frac{\frac{x+c}{2^r}+\frac{d}{2^r}i}{q}\Big)_4=\Big(\frac{2}{q}\Big)\Big(\frac{1+i}{q}\Big)_4\Big(\frac{\frac{x+c+d}{2^{r+1}}-\frac{x+c-d}{2^{r+1}}i}{q}\Big)_4.\end{aligned}\tag{4.3}$$

As $2^r\parallel x+c$ and $2^r\parallel d$ we have $2^{r+1}\mid x+c\pm d$ and $\frac{d+x+c}{2^r}\cdot\frac{d-x-c}{2^r}=(\frac{d}{2^r})^2-(\frac{x+c}{2^r})^2\equiv 0 \pmod 8$. Since $2qy^2=(d+x+c)(d-x-c)+2c(x+c)$ and $r\ge 2$ we see that

$$c\frac{x+c}{2^r}=\frac{qy^2}{2^r}-\frac{(d+x+c)(d-x-c)}{2^{r+1}}\equiv q\frac{y^2}{2^r} \pmod{2^{r+2}}.\tag{4.4}$$

Thus, $r=2t$, $(-1)^{\frac{d}{4}+\frac{y}{2}}=1$ and $\frac{x+c}{2^r}\equiv cq \pmod 8$. Hence

$$\frac{x+c}{2^r}\equiv q\equiv(-1)^{\frac{q-1}{2}}\equiv(-1)^{\frac{q-1}{2}}d_0\equiv(-1)^{\frac{q-1}{2}}\frac{d}{2^r} \pmod 4.$$

Set

$$A=\frac{x+c+(-1)^{\frac{q-1}{2}}d}{2^{r+1}} \quad \text{and} \quad B=\frac{x+c-(-1)^{\frac{q-1}{2}}d}{2^{r+1}}.$$

Then

$$2\nmid A,\ 2\mid B \quad \text{and} \quad A^2+B^2=\frac{(x+c)^2+d^2}{2^{2r+1}}.$$

Thus,

$$\begin{aligned}\Big(\frac{A-Bi}{q}\Big)_4 &= (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}\Big(\frac{q}{A-Bi}\Big)_4 = (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}\Big(\frac{qy^2}{A-Bi}\Big)_4\Big(\frac{y^2}{A-Bi}\Big)_4\\ &= (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}\Big(\frac{((x+c)^2+d^2)/2-x(x+c)}{A-Bi}\Big)_4\Big(\frac{y}{A^2+B^2}\Big)\\ &= (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}\Big(\frac{-2^rx(x+c)/2^r}{A-Bi}\Big)_4\Big(\frac{y}{((x+c)^2+d^2)/2^{2r+1}}\Big)\\ &= (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}\Big(\frac{2}{A-Bi}\Big)_4^r\Big(\frac{-x(x+c)/2^r}{A-Bi}\Big)_4\Big(\frac{2^ty_0}{((x+c)^2+d^2)/2^{2r+1}}\Big)\\ &= (-1)^{\frac{q-1}{2}\cdot\frac{B}{2}}i^{-\frac{AB}{2}r}(-1)^{\frac{-x(x+c)/2^r-1}{2}\cdot\frac{B}{2}}\Big(\frac{A-Bi}{-x(x+c)/2^r}\Big)_4\\ &\quad\times(-1)^{\frac{\frac{1}{2}((\frac{x+c}{2^r})^2+(\frac{d}{2^r})^2)-1}{4}t}\Big(\frac{((x+c)^2+d^2)/2^{2r+1}}{y_0}\Big).\end{aligned}$$

Since $x\equiv 3 \pmod 4$ and $\frac{x+c}{2^r}\equiv q \pmod 4$ we have $(-1)^{\frac{-x(x+c)/2^r-1}{2}}=(-1)^{\frac{q-1}{2}}$. As $\frac{x+c}{2^r}\equiv cq \pmod 8$, we see that $\frac{B}{2}=\frac{1}{4}(\frac{x+c}{2^r}-(-1)^{\frac{q-1}{2}}\frac{d}{2^r})\equiv\frac{cq-(-1)^{\frac{q-1}{2}}d_0}{4} \pmod 2$ and

$$\begin{aligned}(-1)^{\frac{\frac{1}{2}((\frac{x+c}{2^r})^2+(\frac{d}{2^r})^2)-1}{4}} &= (-1)^{\frac{(\frac{x+c}{2^r})^2-1}{8}+\frac{(\frac{d}{2^r})^2-1}{8}} = (-1)^{\frac{(cq)^2-1}{8}+\frac{d_0^2-1}{8}}\\ &= (-1)^{\frac{c^2-1}{8}+\frac{q^2-1}{8}+\frac{d_0^2-1}{8}} = (-1)^{\frac{p-1}{8}+\frac{q^2-1}{8}+\frac{d_0^2-1}{8}}.\end{aligned}$$

Recall that $r=2t$. From the above we deduce that

$$\begin{aligned}&\Big(\frac{A-Bi}{q}\Big)_4\\ &= (-1)^{\frac{AB}{2}t}\cdot(-1)^{(\frac{p-1}{8}+\frac{q^2-1}{8}+\frac{d_0^2-1}{8})t}\Big(\frac{A-Bi}{x(x+c)/2^r}\Big)_4\Big(\frac{((x+c)^2+d^2)/2}{y_0}\Big)\\ &= (-1)^{(\frac{cq-(-1)^{(q-1)/2}d_0}{4}+\frac{p-1}{8}+\frac{q^2-1}{8}+\frac{d_0-1}{4})t}\Big(\frac{A-Bi}{x(x+c)/2^r}\Big)_4\Big(\frac{((x+c)^2+d^2)/2}{y_0}\Big)\\ &= (-1)^{(\frac{cq-(-1)^{(q-1)/2}}{4}+\frac{p-1}{8}+\frac{q^2-1}{8})t}\\ &\quad\times\Big(\frac{x+c+(-1)^{\frac{q-1}{2}}d-(x+c-(-1)^{\frac{q-1}{2}}d)i}{x(x+c)/2^r}\Big)_4\Big(\frac{qy^2+x(x+c)}{y_0}\Big)\\ &= (-1)^{(\frac{c-1}{4}+\frac{q-(-1)^{(q-1)/2}}{4}+\frac{p-1}{8}+\frac{q^2-1}{8})t}\\ &\quad\times\Big(\frac{c+(-1)^{\frac{q-1}{2}}d-(c-(-1)^{\frac{q-1}{2}}d)i}{x}\Big)_4\Big(\frac{(-1)^{\frac{q-1}{2}}d+(-1)^{\frac{q-1}{2}}di}{(x+c)/2^r}\Big)_4\Big(\frac{x(x+c)}{y_0}\Big)\\ &= \Big(\frac{-i(1+i)(c+(-1)^{\frac{q-1}{2}}di)}{x}\Big)_4\Big(\frac{1+i}{(x+c)/2^r}\Big)_4\Big(\frac{x(x+c)}{y_0}\Big).\end{aligned}$$

Recall that $\frac{x+c}{2^r} \equiv cq \pmod 8$. From the above and Lemma 2.7 we deduce that

$$\begin{aligned}\Big(\frac{A-Bi}{q}\Big)_4 &= \Big(\frac{i}{x}\Big)_4\Big(\frac{1+i}{x}\Big)_4\Big(\frac{c+(-1)^{\frac{q-1}{2}}di}{x}\Big)_4 i^{\frac{(-1)^{\frac{(x+c)/2^r-1}{2}}\frac{x+c}{2^r}-1}{4}}\Big(\frac{x(x+c)}{y_0}\Big)\\ &= (-1)^{\frac{x+1}{4}}\cdot i^{\frac{-x-1}{4}}\Big(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\Big)_4 i^{\frac{(-1)^{(q-1)/2}\frac{x+c}{2^r}-1}{4}}\Big(\frac{d+ci}{y_0}\Big)_4\end{aligned}$$

Using (4.4) we see that

$$\begin{aligned}i^{\frac{(-1)^{\frac{q-1}{2}}\frac{x+c}{2^r}-1}{4}} &= i^{\frac{(-1)^{\frac{q-1}{2}}c\frac{x+c}{2^r}-c}{4}} = i^{\frac{(-1)^{\frac{q-1}{2}}qy_0^2-c}{4}} = i^{\frac{(\frac{-1}{q})q(y_0^2-1)+(\frac{-1}{q})-c}{4}}\\ &= (-1)^{\frac{y_0^2-1}{8}} i^{\frac{(\frac{-1}{q})q-c}{4}} = \Big(\frac{i}{y_0}\Big)_4 i^{\frac{(\frac{-1}{q})q-c}{4}}.\end{aligned}$$

Also,

$$i^{-\frac{x+1}{4}} = i^{-\frac{x+c}{4}}\cdot i^{\frac{c-1}{4}} = \begin{cases} i^{\frac{c-1}{4}} & \text{if } r>2,\\ i^{-q+\frac{c-1}{4}} = (-1)^{\frac{q+1}{2}} i^{1+\frac{c-1}{4}} & \text{if } r=2\end{cases}$$

and

$$\begin{aligned}\Big(\frac{i}{y_0}\Big)_4\Big(\frac{d+ci}{y_0}\Big)_4 &= \Big(\frac{c-di}{y_0}\Big)_4 = \Big(\frac{y_0}{c-di}\Big)_4 = \Big(\frac{2}{c-di}\Big)_4^{-t}\Big(\frac{y}{c+di}\Big)_4\\ &= i^{\frac{d}{2}t}\Big(\frac{y}{c-di}\Big)_4 = (-1)^{\frac{d}{4}\cdot\frac{r}{2}}\Big(\frac{y}{c-di}\Big)_4\\ &= (-1)^{\frac{d}{4}}\Big(\frac{y}{c+di}\Big)_4^{-1} = (-1)^{\frac{d}{4}}\Big(\frac{y^{-1}}{c+di}\Big)_4.\end{aligned}$$

Thus,

$$\begin{aligned}\Big(\frac{A-Bi}{q}\Big)_4 &= (-1)^{\frac{x+1}{4}} i^{-\frac{x+1}{4}+\frac{(\frac{-1}{q})q-c}{4}}\Big(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\Big)_4(-1)^{\frac{d}{4}}\Big(\frac{y^{-1}}{c+di}\Big)_4\\ &= (-1)^{\frac{x+1}{4}+\frac{d}{4}} i^{-\frac{x+c}{4}+\frac{(\frac{-1}{q})q-1}{4}}\Big(\frac{x}{c+(-1)^{\frac{q-1}{2}}di}\Big)_4\Big(\frac{y^{-1}}{c+di}\Big)_4.\end{aligned}$$

For $r=2$ we have $\frac{x+c}{4}\equiv q \pmod 4$ and so $i^{-\frac{x+c}{4}} = i^{-q} = (-1)^{\frac{q+1}{2}}i$. For $r>2$ we have $r=2t\ge 4$, $2^4 \mid x+c$ and so $i^{-\frac{x+c}{4}}=1$. If $q\equiv 1 \pmod 4$, from (4.3) and the above we get

$$\begin{aligned}i^k &= \Big(\frac{2}{q}\Big)\Big(\frac{1+i}{q}\Big)_4\Big(\frac{A-Bi}{q}\Big)_4\\ &= (-1)^{\frac{q-1}{4}} i^{\frac{q-1}{4}}(-1)^{\frac{x+1}{4}+\frac{d}{4}} i^{-\frac{x+c}{4}+\frac{q-1}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{y^{-1}}{c+di}\Big)_4\\ &= (-1)^{\frac{x+1}{4}+\frac{d}{4}} i^{-\frac{x+c}{4}}\Big(\frac{x/y}{c+di}\Big)_4 = (-1)^{\frac{x+c}{4}+\frac{d}{4}+\frac{1-c}{4}} i^{-\frac{x+c}{4}}\Big(\frac{x/y}{c+di}\Big)_4\\ &= (-1)^{\frac{1-c}{4}} i^{-\frac{x+c}{4}}\Big(\frac{x/y}{c+di}\Big)_4 = (-1)^{\frac{p-1}{8}} i^{-\frac{x+c}{4}}\Big(\frac{x/y}{c+di}\Big)_4.\end{aligned}$$

and so

$$\Big(\frac{x/y}{c+di}\Big)_4=\begin{cases}(-1)^{\frac{p-1}{8}}i^{k+1} & \text{if } r=2,\\ (-1)^{\frac{p-1}{8}}i^{k} & \text{if } r>2.\end{cases}$$

Now applying Lemma 2.8 we deduce the result. If $q\equiv 3 \pmod 4$, then $(\frac{y}{c^2+d^2})=(\frac{y_0}{p})=(\frac{p}{y_0})=(\frac{x^2+2qy^2}{y_0})=1$. From (4.3) and the above we deduce that

$$\begin{aligned}\Big(\frac{A-Bi}{q}\Big)_4&=(-1)^{\frac{x+1}{4}+\frac{d}{4}}i^{-\frac{x+c}{4}+\frac{-q-1}{4}}\Big(\frac{x}{c-di}\Big)_4\Big(\frac{y^{-1}}{c+di}\Big)_4\\
&=(-1)^{\frac{x+c}{4}+\frac{d}{4}+\frac{1-c}{4}}i^{-\frac{x+c}{4}-\frac{q+1}{4}}\Big(\frac{x}{c+di}\Big)_4^{-1}\Big(\frac{y}{c+di}\Big)_4\Big(\frac{y^{-2}}{c+di}\Big)_4\\
&=(-1)^{\frac{1-c}{4}}i^{-\frac{x+c}{4}-\frac{q+1}{4}}\Big(\frac{x^{-1}y}{c+di}\Big)_4\Big(\frac{y^2}{c+di}\Big)_4^{-1}\\
&=(-1)^{\frac{c^2-1}{8}}i^{-\frac{x+c}{4}-\frac{q+1}{4}}\Big(\frac{x/y}{c+di}\Big)_4^{-1}\Big(\frac{y}{c^2+d^2}\Big)^{-1}\\
&=(-1)^{\frac{p-1}{8}}i^{-\frac{x+c}{4}-\frac{q+1}{4}}\Big(\frac{x/y}{c+di}\Big)_4^{-1}.\end{aligned}$$

Thus, using (4.3) we see that

$$i^k=\Big(\frac{1+i}{q}\Big)_4\Big(\frac{A-Bi}{q}\Big)_4^{-1}=i^{\frac{-q-1}{4}}\cdot(-1)^{\frac{p-1}{8}}i^{\frac{x+c}{4}+\frac{q+1}{4}}\Big(\frac{x/y}{c+di}\Big)_4$$

and so

$$\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{p-1}{8}}i^{k-\frac{x+c}{4}}=\begin{cases}(-1)^{\frac{p-1}{8}}i^{k-q}=(-1)^{\frac{p-1}{8}}i^{k+1} & \text{if } r=2,\\ (-1)^{\frac{p-1}{8}}i^{k} & \text{if } r>2.\end{cases}$$

Applying Lemma 2.8 we deduce the result.

Case 3. $m>r$. Using Lemmas 2.1-2.5 we see that

$$\begin{aligned}i^k&=\Big(\frac{-d/(x+c)+i}{q}\Big)_4=\Big(\frac{\frac{d}{2^r}-\frac{x+c}{2^r}i}{q}\Big)_4=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}\Big(\frac{q}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\\
&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}\Big(\frac{qy^2}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\Big(\frac{y^2}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\\
&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}\Big(\frac{((x+c)^2+d^2)/2-x(x+c)}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\Big(\frac{y}{(\frac{d}{2^r})^2+(\frac{x+c}{2^r})^2}\Big)\\
&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}\Big(\frac{-2^mx(x+c)/2^m}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\Big(\frac{y}{((x+c)^2+d^2)/2^{2r}}\Big)\\
&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}\Big(\frac{2}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4^m\Big(\frac{-x(x+c)/2^m}{\frac{d}{2^r}-\frac{x+c}{2^r}i}\Big)_4\Big(\frac{2^ty_0}{((x+c)^2+d^2)/2^{2r}}\Big)\\
&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}i^{-\frac{d}{2^r}\cdot\frac{x+c}{2^{r+1}}\cdot m}(-1)^{\frac{-x(x+c)/2^m-1}{2}\cdot\frac{x+c}{2^{r+1}}}\\
&\quad\times\Big(\frac{\frac{d}{2^r}-\frac{x+c}{2^r}i}{-x(x+c)/2^m}\Big)_4(-1)^{\frac{x+c}{2^{r+1}}t}\Big(\frac{(x+c)^2+d^2}{y_0}\Big).\end{aligned}$$

Applying Lemma 2.7 we derive that

$$\begin{aligned}
&\Big(\frac{\frac{d}{2^r}-\frac{x+c}{2^r}i}{-x(x+c)/2^m}\Big)_4\Big(\frac{(x+c)^2+d^2}{y_0}\Big)\\
&=\Big(\frac{d-(x+c)i}{-x}\Big)_4\Big(\frac{d-(x+c)i}{(x+c)/2^m}\Big)_4\Big(\frac{2qy^2+2x(c+x)}{y_0}\Big)\\
&=\Big(\frac{d-ci}{-x}\Big)_4\Big(\frac{d}{(x+c)/2^m}\Big)_4\Big(\frac{2x(c+x)}{y_0}\Big)\\
&=\Big(\frac{-i}{x}\Big)_4\Big(\frac{c+di}{x}\Big)_4\Big(\frac{2}{y_0}\Big)\Big(\frac{d+ci}{y_0}\Big)_4=(-1)^{\frac{x+1}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{i(d+ci)}{y_0}\Big)_4\\
&=(-1)^{\frac{x+1}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{c-di}{y_0}\Big)_4=(-1)^{\frac{x+1}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{y_0}{c-di}\Big)_4\\
&=(-1)^{\frac{x+1}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{y_0}{c+di}\Big)_4^{-1}=(-1)^{\frac{x+c}{4}+\frac{1-c}{4}}\Big(\frac{x}{c+di}\Big)_4\Big(\frac{2^t/y}{c+di}\Big)_4\\
&=(-1)^{\frac{c-1}{4}+\frac{d}{4}t}\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{p-1}{8}+\frac{d}{4}t}\Big(\frac{x/y}{c+di}\Big)_4.
\end{aligned}$$

Therefore,

$$\begin{aligned}
i^k&=(-1)^{\frac{q-1}{2}\cdot\frac{x+c}{2^{r+1}}}i^{-\frac{x+c}{2^{r+1}}\cdot m}(-1)^{\frac{x(x+c)/2^m+1}{2}\cdot\frac{x+c}{2^{r+1}}}\cdot(-1)^{\frac{x+c}{2^{r+1}}t+\frac{p-1}{8}+\frac{d}{4}t}\Big(\frac{x/y}{c+di}\Big)_4\\
&=\begin{cases}(-1)^{\frac{q-1}{2}+\frac{x(x+c)/2^{r+1}+1}{2}+\frac{p-1}{8}+(1+\frac{d}{4})t}i^{-\frac{x+c}{2^{r+1}}(r+1)}\big(\frac{x/y}{c+di}\big)_4 & \text{if } m=r+1,\\ (-1)^{m\frac{x+c}{2^{r+2}}+\frac{p-1}{8}+\frac{d}{4}t}\big(\frac{x/y}{c+di}\big)_4 & \text{if } m>r+1.\end{cases}
\end{aligned}$$

Since $2^t\parallel y$, $2^r\parallel d$, $2^m\parallel x+c$, $m>r\ge 2$ and $2qy^2-d^2=(x+c)^2-2x(x+c)$, we see that $2^{m+1}\parallel 2qy^2-d^2$. That is, $2^{m+1}\parallel 2^{2t+1}qy_0^2-2^{2r}d_0^2$. Hence $m+1=\min\{2t+1,2r\}\le 2r$ and so $t>1$. If $m+1=2r$, then $t\ge r$ and

$$\begin{aligned}
x\frac{x+c}{2^m}&=2^{m-1}\Big(\frac{x+c}{2^m}\Big)^2-2^{2t+1-2r}qy_0^2+d_0^2\\
&\equiv 1-2^{2t+1-2r}q\equiv\begin{cases}1\pmod 4 & \text{if } t>r,\\ 3\pmod 4 & \text{if } t=r.\end{cases}
\end{aligned}$$

If $m+1<2r$, then $m=2t$, $t<r$ and

$$\begin{aligned}
x\frac{x+c}{2^m}&=2^{m-1}\Big(\frac{x+c}{2^m}\Big)^2-qy_0^2+2^{2r-1-2t}d_0^2\\
&\equiv -q+2^{2r-2t-1}\equiv\begin{cases}-q\pmod 4 & \text{if } t<r-1,\\ 2-q\pmod 4 & \text{if } t=r-1.\end{cases}
\end{aligned}$$

For $r=2$ we must have $m=3$ and so

$$\begin{aligned}
i^k&=(-1)^{\frac{q-1}{2}+\frac{x(x+c)/8+1}{2}+\frac{p-1}{8}+(1+\frac{d}{4})t}i^{-\frac{x+c}{8}\cdot 3}\Big(\frac{x/y}{c+di}\Big)_4\\
&=(-1)^{\frac{q-1}{2}+\frac{x(x+c)/8+1}{2}+\frac{p-1}{8}}i^{\frac{x+c}{8}}\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{q-1}{2}+\frac{p-1}{8}}i\Big(\frac{x/y}{c+di}\Big)_4.
\end{aligned}$$

Now assume $r>2$. If $m+1=2r$, then $m>r+1$ and so

$$i^k=(-1)^{\frac{x+c}{2^{r+2}}+\frac{p-1}{8}}\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{2^{r-3}+\frac{p-1}{8}}\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{d}{8}+\frac{p-1}{8}}\Big(\frac{x/y}{c+di}\Big)_4.$$

If $m+1<2r$ and $m=r+1$, then $t=\frac{m}{2}=\frac{r+1}{2}$, $x\frac{x+c}{2^{r+1}}\equiv -q+2^{2r-2t-1}=-q+2^{r-2}\equiv -q+\frac{d}{4} \pmod 4$ and so

$$\begin{aligned}i^k&=(-1)^{\frac{q-1}{2}+\frac{x(x+c)/2^{r+1}+1}{2}+\frac{p-1}{8}+t}i^{-\frac{x+c}{2^{r+1}}(r+1)}\Big(\frac{x/y}{c+di}\Big)_4\\&=(-1)^{\frac{q-1}{2}+\frac{1-q+\frac{d}{4}}{2}+\frac{p-1}{8}+\frac{r+1}{2}}(-1)^{\frac{r+1}{2}}\Big(\frac{x/y}{c+di}\Big)_4=(-1)^{\frac{p-1}{8}+\frac{d}{8}}\Big(\frac{x/y}{c+di}\Big)_4.\end{aligned}$$

If $m+1<2r$ and $m>r+1$, then $r>3$, $m=2t$ and so $i^k=(-1)^{\frac{p-1}{8}}\big(\frac{x/y}{c+di}\big)_4=(-1)^{\frac{p-1}{8}+\frac{d}{8}}\big(\frac{x/y}{c+di}\big)_4$. Hence

$$\Big(\frac{x/y}{c+di}\Big)_4=\begin{cases}(-1)^{\frac{q-1}{2}+\frac{p-1}{8}}i^{k-1} & \text{if } r=2,\\ (-1)^{\frac{p-1}{8}+\frac{d}{8}}i^k & \text{if } r>2.\end{cases}$$

Now applying Lemma 2.8 we obtain the result.

By the above, the theorem is proved.

Theorem 4.3. *Let p and q be primes such that $p\equiv 1 \pmod 8$ and $q\equiv 3 \pmod 4$. Suppose $p=c^2+d^2=x^2+2qy^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1 \pmod 4$, $d=2^rd_0$, $d_0\equiv 1 \pmod 4$ and $(\frac{c-di}{x})^{\frac{q+1}{4}}\equiv i^m \pmod q$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then*

$$(-q)^{\frac{p-1}{8}}\equiv(-1)^{\frac{x-1}{2}\cdot\frac{q+1}{4}}\Big(\frac{d}{c}\Big)^m \pmod p.$$

Proof. Clearly $q\nmid x$ and x is odd. We first assume $(c,x+d)=1$. By the proof of Theorem 4.1, $(q,(x+d)(c^2+(x+d)^2))=1$. It is easily seen that $\frac{c/(x+d)-i}{c/(x+d)+i}=\frac{c-(x+d)i}{c+(x+d)i}\equiv\frac{c-di}{ix} \pmod q$. Thus, for $k=0,1,2,3$, using Lemma 2.6 we get

$$\begin{aligned}&\Big(\frac{c/(x+d)+i}{q}\Big)_4=i^k\\&\iff \frac{c}{x+d}\in Q_k(q)\iff\Big(\frac{\frac{c}{x+d}-i}{\frac{c}{x+d}+i}\Big)^{\frac{q+1}{4}}\equiv i^k \pmod q\\&\iff\Big(\frac{c-di}{ix}\Big)^{\frac{q+1}{4}}\equiv i^k \pmod q\iff\Big(\frac{c-di}{x}\Big)^{\frac{q+1}{4}}\equiv i^{\frac{q+1}{4}+k} \pmod q.\end{aligned}$$

Since $(\frac{c-di}{x})^{\frac{q+1}{4}}\equiv i^m \pmod q$, from the above we deduce that

$$\Big(\frac{c/(x+d)+i}{q}\Big)_4=i^{m-\frac{q+1}{4}}=\begin{cases}(-1)^{\frac{q+5}{8}}i^{m+1} & \text{if } q\equiv 3 \pmod 8,\\ (-1)^{\frac{q+1}{8}}i^m & \text{if } q\equiv 7 \pmod 8.\end{cases}$$

Now, applying Theorem 4.1 we derive the result.

Now we assume $(d_0, x+c) = 1$. By the proof of Theorem 4.2, $(q, x+c) = (q, d^2+(x+c)^2) = 1$. It is easily seen that $\frac{d+(x+c)i}{d-(x+c)i} \equiv \frac{c-di}{-x} \pmod q$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\begin{aligned}
&\Big(\frac{-d/(x+c)+i}{q}\Big)_4 = i^k \\
&\iff -\frac{d}{x+c} \in Q_k(q) \iff \Big(\frac{-\frac{d}{x+c}-i}{-\frac{d}{x+c}+i}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \\
&\iff \Big(\frac{d+(x+c)i}{d-(x+c)i}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \iff \Big(\frac{c-di}{-x}\Big)^{\frac{q+1}{4}} \equiv i^k \pmod q \\
&\iff \Big(\frac{c-di}{x}\Big)^{\frac{q+1}{4}} \equiv i^{\frac{q+1}{2}+k} \pmod q.
\end{aligned}$$

Since $(\frac{c-di}{x})^{\frac{q+1}{4}} \equiv i^m \pmod q$, from the above we deduce that $(\frac{-d/(x+c)+i}{q})_4 = i^{m-\frac{q+1}{2}} = (-1)^{\frac{q+1}{4}} i^m$. Now applying Theorem 4.2 we deduce the result. The proof is now complete.

Corollary 4.1. *Let p and q be primes such that $p \equiv 1 \pmod 8$ and $q \equiv 3 \pmod 8$. Suppose $p = c^2+d^2 = x^2+2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$, $q \mid cd$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{x-1}{2}} \pmod p & \text{if } x \equiv \pm c \pmod q, \\ \mp(-1)^{\frac{q-3}{8}+\frac{x-1}{2}} \frac{d}{c} \pmod p & \text{if } x \equiv \pm d \pmod q. \end{cases}$$

Proof. If $x \equiv \pm c \pmod q$, then $q \mid d$ and so $(\frac{c-di}{x})^{\frac{q+1}{4}} \equiv (\pm 1)^{\frac{q+1}{4}} = \pm 1 \pmod q$. If $x \equiv \pm d \pmod q$, then $q \mid c$ and so $(\frac{c-di}{x})^{\frac{q+1}{4}} \equiv (\mp i)^{\frac{q+1}{4}} = \mp(-1)^{\frac{q-3}{8}} i \pmod q$. Now applying Theorem 4.3 we deduce the result.

As an example, taking $q = 3$ in Corollary 4.1 we see that if p is a prime of the form $24k+1$ and so $p = c^2+d^2 = x^2+6y^2$, then

$$(-3)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{x-1}{2}} \pmod p & \text{if } x \equiv \pm c \pmod 3, \\ \mp(-1)^{\frac{x-1}{2}} \frac{d}{c} \pmod p & \text{if } x \equiv \pm d \pmod 3. \end{cases} \tag{4.5}$$

Theorem 4.4. *Let p and q be primes such that $p \equiv 1 \pmod 8$, $q \equiv 7 \pmod 8$, $p = c^2+d^2 = x^2+2qy^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \Big(\frac{d}{c}\Big)^m \pmod p \iff \Big(\frac{c-di}{c+di}\Big)^{\frac{q+1}{8}} \equiv i^m \pmod q.$$

Proof. Observe that

$$\Big(\frac{c-di}{c+di}\Big)^{\frac{q+1}{8}} = \frac{(c-di)^{\frac{q+1}{4}}}{(c^2+d^2)^{\frac{q+1}{8}}} = \frac{(c-di)^{\frac{q+1}{4}}}{(x^2+2qy^2)^{\frac{q+1}{8}}} \equiv \Big(\frac{c-di}{x}\Big)^{\frac{q+1}{4}} \pmod q.$$

The result follows from Theorem 4.3.

We note that if $q \nmid d$, then the m in Theorem 4.4 depends only on $\frac{c}{d} \pmod q$.

Corollary 4.2. *Let $p \equiv 1 \pmod 8$ and $q \equiv 7 \pmod 8$ be primes such that $p = c^2+d^2 = x^2+2qy^2$ with $c,d,x,y \in \mathbb{Z}$ and $q \mid cd(c^2-d^2)$. Suppose $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$q^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q+1}{8}} \pmod p & \text{if } q \mid c, \\ 1 \pmod p & \text{if } q \mid d, \\ \pm(-1)^{\frac{q+9}{16}}\frac{d}{c} \pmod p & \text{if } 16 \mid (q-7) \text{ and } c \equiv \pm d \pmod q, \\ (-1)^{\frac{q+1}{16}} \pmod p & \text{if } 16 \mid (q-15) \text{ and } c \equiv \pm d \pmod q. \end{cases}$$

Proof. Clearly

$$\frac{c-di}{c+di} \equiv \begin{cases} -1 \pmod q & \text{if } q \mid c, \\ 1 \pmod q & \text{if } q \mid d, \\ -i \pmod q & \text{if } c \equiv d \pmod q, \\ i \pmod q & \text{if } c \equiv -d \pmod q. \end{cases}$$

Thus the result follows from Theorem 4.4.

Theorem 4.5. *Let p and q be distinct primes, $p \equiv 1 \pmod 8$, $q \equiv 1 \pmod 4$, $p = c^2+d^2 = x^2+2qy^2$, $q = a^2+b^2$, $a,b,c,d,x,y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Suppose $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{b}{a})^m \pmod q$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{x-1}{2}\cdot\frac{q-1}{4}+\frac{d}{4}+\frac{y}{2}}\Big(\frac{d}{c}\Big)^m \pmod p.$$

Proof. Clearly $q \nmid x$. We first assume $(c, x+d) = 1$. By the proof of Theorem 4.1, $(q, (x+d)(c^2+(x+d)^2)) = 1$. It is easily seen that $\frac{ac+b(x+d)}{ac-b(x+d)} \equiv \frac{ac+bd}{ax}\cdot\frac{b}{a} \pmod q$. Thus, for $k = 0,1,2,3$, using Lemma 2.6 we get

$$\begin{aligned} \Big(\frac{c/(x+d)+i}{q}\Big)_4 = i^k &\iff \frac{c}{x+d} \in Q_k(q) \iff \Big(\frac{\frac{c}{x+d}+\frac{b}{a}}{\frac{c}{x+d}-\frac{b}{a}}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q \\ &\iff \Big(\frac{ac+b(x+d)}{ac-b(x+d)}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q \\ &\iff \Big(\frac{ac+bd}{ax}\cdot\frac{b}{a}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q \\ &\iff \Big(\frac{ac+bd}{ax}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^{k-\frac{q-1}{4}} \pmod q. \end{aligned}$$

Since $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{b}{a})^m \pmod q$, from the above we get $(\frac{c/(x+d)+i}{q})_4 = i^{m+\frac{q-1}{4}}$. Now the result follows from Theorem 4.1 immediately.

Suppose $(d_0, x+c) = 1$. By the proof of Theorem 4.2, $(q, (x+c)(d^2+(x+c)^2)) = 1$. It is easily seen that $\frac{ad-b(x+c)}{ad+b(x+c)} \equiv \frac{ac+bd}{-ax} \pmod q$. Thus, for $k = 0, 1, 2, 3$, using Lemma 2.6 we get

$$\Big(\frac{-d/(x+c)+i}{q}\Big)_4 = i^k$$

$$\iff -\frac{d}{x+c} \in Q_k(q) \iff \Big(\frac{-\frac{d}{x+c}+\frac{b}{a}}{-\frac{d}{x+c}-\frac{b}{a}}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q$$

$$\iff \Big(\frac{ad-b(x+c)}{ad+b(x+c)}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q \iff \Big(\frac{ac+bd}{-ax}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^k \pmod q$$

$$\iff \Big(\frac{ac+bd}{ax}\Big)^{\frac{q-1}{4}} \equiv \Big(\frac{b}{a}\Big)^{\frac{q-1}{2}+k} \pmod q.$$

Since $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{b}{a})^m \pmod q$, by the above we get $(\frac{-d/(x+c)+i}{q})_4 = i^{m-\frac{q-1}{2}}$. Now applying Theorem 4.2 we derive the result. The proof is now complete.

Corollary 4.3. *Let $p \equiv 1 \pmod 8$ and $q \equiv 5 \pmod 8$ be primes such that $p = c^2+d^2 = x^2+2qy^2$ with $c, d, x, y \in \mathbb{Z}$ and $q \mid cd$. Suppose $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}} \pmod p & \text{if } x \equiv \pm c \pmod q, \\ \pm(-1)^{\frac{q-5}{8}+\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}\frac{d}{c}} \pmod p & \text{if } x \equiv \pm d \pmod q. \end{cases}$$

Proof. Suppose $q = a^2+b^2$ with $a, b \in \mathbb{Z}$. If $x \equiv \pm c \pmod q$, then $q \mid d$ and so $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{c}{x})^{\frac{q-1}{4}} \equiv (\pm 1)^{\frac{q-1}{4}} = \pm 1 \pmod q$. If $x \equiv \pm d \pmod q$, then $q \mid c$ and so $(\frac{ac+bd}{ax})^{\frac{q-1}{4}} \equiv (\frac{bd}{ax})^{\frac{q-1}{4}} \equiv (\pm\frac{b}{a})^{\frac{q-1}{4}} \equiv \pm(-1)^{\frac{q-5}{8}}\frac{b}{a} \pmod q$. Now putting the above with Theorem 4.5 we deduce the result.

Theorem 4.6. *Let p and q be distinct primes such that $p \equiv 1 \pmod 8$, $q \equiv 1 \pmod 8$, $p = c^2+d^2 = x^2+2qy^2$, $q = a^2+b^2$, $a, b, c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv (-1)^{\frac{d}{4}+\frac{y}{2}}\Big(\frac{d}{c}\Big)^m \pmod p \iff \Big(\frac{ac+bd}{ac-bd}\Big)^{\frac{q-1}{8}} \equiv \Big(\frac{b}{a}\Big)^m \pmod q.$$

Proof. Observe that $b^2 \equiv -a^2 \pmod q$, $p \equiv x^2 \pmod q$ and so

$$\Big(\frac{ac+bd}{ac-bd}\Big)^{\frac{q-1}{8}} = \frac{(ac+bd)^{\frac{q-1}{4}}}{(a^2c^2-b^2d^2)^{\frac{q-1}{8}}} \equiv \frac{(ac+bd)^{\frac{q-1}{4}}}{(a^2p)^{\frac{q-1}{8}}} \equiv \Big(\frac{ac+bd}{ax}\Big)^{\frac{q-1}{4}} \pmod q.$$

The result follows from Theorem 4.5.

We note that if $q \nmid d$, then the m in Theorem 4.6 depends only on $\frac{c}{d} \pmod q$.

Corollary 4.4. *Let p and q be distinct primes of the form $8k+1$ such that $p = c^2+d^2 = x^2+2qy^2$ with $c,d,x,y \in \mathbb{Z}$ and $q \mid cd(c^2-d^2)$. Suppose $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-q)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8}+\frac{d}{4}+\frac{y}{2}} \pmod p & \text{if } q \mid c, \\ (-1)^{\frac{d}{4}+\frac{y}{2}} \pmod p & \text{if } q \mid d, \\ (-1)^{\frac{q-1}{16}+\frac{d}{4}+\frac{y}{2}} \pmod p & \text{if } 16 \mid (q-1) \text{ and } c \equiv \pm d \pmod q, \\ \pm(-1)^{\frac{q-9}{16}+\frac{d}{4}+\frac{y}{2}}\frac{d}{c} \pmod p & \text{if } 16 \mid (q-9) \text{ and } c \equiv \pm d \pmod q. \end{cases}$$

Proof. Suppose that $q = a^2+b^2$ with $a, b \in \mathbb{Z}$. Then clearly

$$\Big(\frac{ac+bd}{ac-bd}\Big)^{\frac{q-1}{8}} \equiv \begin{cases} (-1)^{\frac{q-1}{8}} \pmod q & \text{if } q \mid c, \\ 1 \pmod q & \text{if } q \mid d, \\ (-1)^{\frac{q-1}{16}} \pmod q & \text{if } 16 \mid (q-1) \text{ and } c \equiv \pm d \pmod q, \\ \pm(-1)^{\frac{q-9}{16}}\frac{b}{a} \pmod q & \text{if } 16 \mid (q-9) \text{ and } c \equiv \pm d \pmod q. \end{cases}$$

Thus the result follows from Theorem 4.6.

Theorem 4.7. *Let $p \equiv 1 \pmod 8$ be a prime, $p = c^2+d^2 = x^2+2(a^2+b^2)y^2$, $a,b,c,d,x,y \in \mathbb{Z}$, $a \neq 0$, $2 \mid a$, $(a,b) = 1$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x+d) = 1$ or $(d_0, x+c) = 1$. Then*

$$(-a^2-b^2)^{\frac{p-1}{8}} \equiv \begin{cases} (-1)^{\frac{d}{4}+\frac{y}{2}}(\frac{c}{d})^m \pmod p & \text{if } 4 \mid a, \\ (-1)^{\frac{b-1}{2}+\frac{d}{4}+\frac{y}{2}+\frac{x-1}{2}}(\frac{c}{d})^{m-1} \pmod p & \text{if } 4 \mid a-2. \end{cases}$$

$$\iff \Big(\frac{(ac+bd)/x}{b+ai}\Big)_4 = i^m.$$

Proof. Suppose $q = a^2+b^2$ and $(\frac{(ac+bd)/x}{b+ai})_4 = i^m$. Then clearly $q \equiv 1 \pmod 4$ and $p \nmid q$. We first assume $(c, x+d) = 1$. By the proof of Theorem 4.1, $(q, x+d) = (q, c^2+(x+d)^2) = 1$. Since $\frac{c-(x+d)i}{c+(x+d)i} \equiv \frac{c-di}{ix} \pmod q$, from [S5, p.24] we know that

$$\Big(\frac{c/(x+d)+i}{q}\Big)_4 = (-1)^{\frac{b+1}{2}\cdot\frac{a}{2}+[\frac{q}{8}]}i^{-m}.$$

This together with Theorem 4.1 yields the result in this case.

Now we assume $(d_0, x+c) = 1$. By the proof of Theorem 4.2, $(q, x+c) = (q, (x+c)^2+d^2) = 1$. Since $\frac{d+(x+c)i}{d-(x+c)i} \equiv \frac{c-di}{-x} \pmod q$, from [S5, p.24] we know that

$$\Big(\frac{-d/(x+c)+i}{q}\Big)_4 = \begin{cases} (-1)^{\frac{b+1}{2}}i^{1-m} & \text{if } 4 \mid a-2, \\ i^{-m} & \text{if } 4 \mid a. \end{cases}$$

Now applying Theorem 4.2 we deduce the result in this case. So the theorem is proved.

Corollary 4.5. *Let $p \equiv 1, 9 \pmod{40}$ be a prime and so $p = c^2 + d^2 = x^2 + 10y^2$ with $c, d, x, y \in \mathbb{Z}$. Suppose $c \equiv 1 \pmod 4$, $d = 2^r d_0$ and $d_0 \equiv 1 \pmod 4$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$(-5)^{\frac{p-1}{8}} \equiv \begin{cases} \pm(-1)^{\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}} \pmod p & \text{if } \frac{2c-d}{x} \equiv \pm 2 \pmod 5, \\ \pm(-1)^{\frac{d}{4}+\frac{x-1}{2}+\frac{y}{2}} \frac{c}{d} \pmod p & \text{if } \frac{2c-d}{x} \equiv \pm 1 \pmod 5. \end{cases}$$

Proof. Since $(\frac{\pm 1}{1+2i})_4 = \pm 1$ and $(\frac{\pm 2}{1+2i})_4 = \pm i$, taking $a = 2$ and $b = 1$ in Theorem 4.7 we derive the result.

We remark that Corollary 4.5 partially solves [S4, Conjecture 9.8].

5. Congruences for $U_{\frac{p-1}{4}}(2a, -1)$ and $V_{\frac{p-1}{4}}(2a, -1) \pmod p$.

For two numbers P and Q the Lucas sequences $\{U_n(P, Q)\}$ and $\{V_n(P, Q)\}$ are defined by

$$U_0(P, Q) = 0,\ U_1(P, Q) = 1,\ U_{n+1}(P, Q) = PU_n(P, Q) - QU_{n-1}(P, Q)\ (n \ge 1),$$
$$V_0(P, Q) = 2,\ V_1(P, Q) = P,\ V_{n+1}(P, Q) = PV_n(P, Q) - QV_{n-1}(P, Q)\ (n \ge 1).$$

Set $D = P^2 - 4Q$. It is well known that

$$U_n(P, Q) = \frac{1}{\sqrt{D}}\Big\{\Big(\frac{P + \sqrt{D}}{2}\Big)^n - \Big(\frac{P - \sqrt{D}}{2}\Big)^n\Big\} \quad (D \ne 0) \tag{5.1}$$

and

$$V_n(P, Q) = \Big(\frac{P + \sqrt{D}}{2}\Big)^n + \Big(\frac{P - \sqrt{D}}{2}\Big)^n. \tag{5.2}$$

Theorem 5.1. *Let p be a prime of the form $8k + 1$ and $a \in \mathbb{Z}$ with $2 \nmid a$. Suppose that $p = c^2 + d^2 = x^2 + (a^2 + 1)y^2$, $c, d, x, y \in \mathbb{Z}$, $c \equiv 1 \pmod 4$, $d = 2^r d_0$, $y = 2^\beta y_0$ and $d_0 \equiv y_0 \equiv 1 \pmod 4$. Assume $(c, x + d) = 1$ or $(d_0, x + c) = 1$. Then*

$$U_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} (-1)^{\frac{a-1}{2}+\frac{d}{4}+\frac{x-1}{2}} \frac{y}{x} \pmod p & \text{if } 4 \mid y - 2, \\ 0 \pmod p & \text{if } 4 \mid y \end{cases}$$

and

$$V_{\frac{p-1}{4}}(2a, -1) \equiv \begin{cases} 0 \pmod p & \text{if } 4 \mid y - 2, \\ 2(-1)^{\frac{d}{4}+\frac{y}{4}} \pmod p & \text{if } 4 \mid y. \end{cases}$$

Proof. Set $a_1 = (1 - (-1)^{\frac{a-1}{2}}a)/2$ and $b_1 = (1 + (-1)^{\frac{a-1}{2}}a)/2$. Then $2 \mid a_1$, $2 \nmid b_1$ and $a^2 + 1 = 2(a_1^2 + b_1^2)$. It is clear that

$$\Big(\frac{(a_1c + b_1d)/((-1)^{\frac{x-1}{2}}x)}{b_1 + a_1 i}\Big)_4 = (-1)^{\frac{x-1}{2}\cdot\frac{a_1}{2}}\Big(\frac{(a_1c + b_1d)/x}{b_1 + a_1 i}\Big)_4.$$

We first assume $a\equiv 1 \pmod 4$. Replacing d, x with $-d, (-1)^{\frac{x-1}{2}}x$ in [S4, Theorem 8.3(i)] we obtain

$$U_{\frac{p-1}{4}}(2a,-1)\equiv\begin{cases}\mp(-1)^{\frac{x-1}{2}\cdot\frac{a-1}{4}}(-a_1^2-b_1^2)^{\frac{p-1}{8}}(-\frac{c}{d})^{(1-(-1)^{\frac{a-1}{4}})/2}(-1)^{\frac{x-1}{2}}\frac{y}{x} \pmod p\\ \qquad\qquad \text{if } 4\mid y-2 \text{ and } \big(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\big)_4=\pm 1,\\ \mp(-1)^{\frac{x-1}{2}\cdot\frac{a-1}{4}}(-a_1^2-b_1^2)^{\frac{p-1}{8}}(-\frac{c}{d})^{1+(1-(-1)^{\frac{a-1}{4}})/2}(-1)^{\frac{x-1}{2}}\frac{y}{x} \pmod p\\ \qquad\qquad \text{if } 4\mid y-2 \text{ and } \big(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\big)_4=\pm i,\\ 0 \pmod p \ \text{ if } 4\mid y\end{cases}$$

and

$$V_{\frac{p-1}{4}}(2a,-1)\equiv\begin{cases}\pm 2(-1)^{\frac{x-1}{2}\cdot\frac{a-1}{4}+\frac{y}{4}}(-a_1^2-b_1^2)^{\frac{p-1}{8}}(-\frac{c}{d})^{(1-(-1)^{\frac{a-1}{4}})/2} \pmod p\\ \qquad\qquad \text{if } 4\mid y \text{ and } \big(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\big)_4=\pm 1,\\ \pm 2(-1)^{\frac{x-1}{2}\cdot\frac{a-1}{4}+\frac{y}{4}}(-a_1^2-b_1^2)^{\frac{p-1}{8}}(-\frac{c}{d})^{1+(1-(-1)^{\frac{a-1}{4}})/2} \pmod p\\ \qquad\qquad \text{if } 4\mid y \text{ and } \big(\frac{(a_1c+b_1d)/x}{b_1+a_1i}\big)_4=\pm i,\\ 0 \pmod p \ \text{ if } 4\mid y-2.\end{cases}$$

From Theorem 4.7 we know that

$$\begin{aligned}&(-a_1^2-b_1^2)^{\frac{p-1}{8}}\\ &\equiv\begin{cases}\pm(-1)^{\frac{d}{4}+\frac{y}{2}} \pmod p & \text{if } 4\mid a_1 \text{ and } (\frac{(a_1c+b_1d)/x}{b_1+a_1i})_4=\pm 1,\\ \pm(-1)^{\frac{b_1-1}{2}+\frac{d}{4}+\frac{y}{2}+\frac{x-1}{2}}\frac{d}{c} \pmod p & \text{if } 4\mid a_1-2 \text{ and } (\frac{(a_1c+b_1d)/x}{b_1+a_1i})_4=\pm 1,\\ \pm(-1)^{\frac{d}{4}+\frac{y}{2}}\frac{c}{d} \pmod p & \text{if } 4\mid a_1 \text{ and } (\frac{(a_1c+b_1d)/x}{b_1+a_1i})_4=\pm i,\\ \pm(-1)^{\frac{b_1-1}{2}+\frac{d}{4}+\frac{y}{2}+\frac{x-1}{2}} \pmod p & \text{if } 4\mid a_1-2 \text{ and } (\frac{(a_1c+b_1d)/x}{b_1+a_1i})_4=\pm i.\end{cases}\end{aligned}$$

Now putting the above together we deduce the result in the case $a\equiv 1 \pmod 4$. The case $a\equiv 3 \pmod 4$ can be proved similarly by using [S4, Theorem 8.3(ii)] and Theorem 4.7.

Corollary 5.1. *Let p be a prime of the form $8k+1$ and $a\in\mathbb{Z}$ with $2\nmid a$. Suppose that $p=c^2+d^2=x^2+(a^2+1)y^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1 \pmod 4$, $d=2^rd_0$, $d_0\equiv 1 \pmod 4$ and $4\mid y$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then*

$$(a+\sqrt{a^2+1})^{\frac{p-1}{4}}\equiv(-1)^{\frac{d}{4}+\frac{y}{4}} \pmod p.$$

Proof. From (5.1), (5.2) and Theorem 5.1 we see that

$$\begin{aligned}(a+\sqrt{a^2+1})^{\frac{p-1}{4}}&=\frac{1}{2}V_{\frac{p-1}{4}}(2a,-1)+\sqrt{a^2+1}U_{\frac{p-1}{4}}(2a,-1)\\ &\equiv\frac{1}{2}V_{\frac{p-1}{4}}(2a,-1)\equiv(-1)^{\frac{d}{4}+\frac{y}{4}} \pmod p.\end{aligned}$$

This proves the corollary.

Corollary 5.2. *Let p be a prime of the form $8k+1$ and $a\in\mathbb{Z}$ with $2\nmid a$. Suppose that $p=c^2+d^2=x^2+(a^2+1)y^2$, $c,d,x,y\in\mathbb{Z}$, $c\equiv 1\pmod 4$, $d=2^rd_0$ and $d_0\equiv 1\pmod 4$. Assume $(c,x+d)=1$ or $(d_0,x+c)=1$. Then*

$$p\mid U_{\frac{p-1}{8}}(2a,-1)\iff 4\mid y\quad\text{and}\quad\frac{p-1}{8}\equiv\frac{d}{4}+\frac{y}{4}\pmod 2.$$

Proof. By [S4, (1.5)],

$$p\mid U_{\frac{p-1}{8}}(2a,-1)\iff V_{\frac{p-1}{4}}(2a,-1)\equiv 2(-1)^{\frac{p-1}{8}}\pmod p.$$

Now applying Theorem 5.1 we obtain the result.

Remark 5.1 Theorem 5.1 and Corollary 5.2 were conjectured by the author in [S4, Conjectures 9.17 and 9.19].

References

[BEW] B.C. Berndt, R.J. Evans and K.S. Williams, *Gauss and Jacobi Sums*, Wiley, New York, 1998.

[E] R.J. Evans, *Residuacity of primes*, Rocky Mountain J. Math. **19** (1989), 1069-1081.

[HW] R.H. Hudson and K.S. Williams, *Extensions of theorems of Cunningham-Aigner and Hasse-Evans*, Pacific J. Math. **104** (1983), 111-132.

[IR] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory, 2nd ed.*, Springer, New York, 1990.

[L] E. Lehmer, *On Euler's criterion*, J. Austral. Math. Soc. **1** (1959), 64-70.

[Lem] F. Lemmermeyer, *Reciprocity Laws: From Euler to Eisenstein*, Springer, Berlin, 2000.

[S1] Z.H. Sun, *Supplements to the theory of quartic residues*, Acta Arith. **97** (2001), 361-377.

[S2] Z.H. Sun, *Quartic residues and binary quadratic forms*, J. Number Theory **113** (2005), 10-52.

[S3] Z.H. Sun, *On the quadratic character of quadratic units*, J. Number Theory **128** (2008), 1295-1335.

[S4] Z.H. Sun, *Quartic, octic residues and Lucas sequences*, J. Number Theory **129** (2009), 499-550.

[S5] Z.H. Sun, *Congruences for $q^{[p/8]}\pmod p$*, Acta Arith. **159** (2013), 1-25.